

Convergence of Sequences of Random Variables¹

- Definitions (All quantities in boldface are vectors in \mathbb{R}^m unless otherwise stated)
 - ★ $\mathbf{X}_n \xrightarrow{a.s.} \mathbf{X}$ means $P\{s : \lim_{n \rightarrow \infty} \mathbf{X}(s) = \mathbf{X}(s)\} = 1$.
 - ★ $\mathbf{X}_n \xrightarrow{P} \mathbf{X}$ means $\forall \epsilon > 0, \lim_{n \rightarrow \infty} P\{|\mathbf{X}_n - \mathbf{X}| < \epsilon\} = 1$.
 - ★ $\mathbf{X}_n \xrightarrow{d} \mathbf{X}$ means for every continuity point \mathbf{x} of $F_{\mathbf{X}}$, $\lim_{n \rightarrow \infty} F_{\mathbf{X}_n}(\mathbf{x}) = F_{\mathbf{X}}(\mathbf{x})$.
- $\mathbf{X}_n \xrightarrow{a.s.} \mathbf{X} \Rightarrow \mathbf{X}_n \xrightarrow{P} \mathbf{X} \Rightarrow \mathbf{X}_n \xrightarrow{d} \mathbf{X}$.
- If \mathbf{a} is a vector of constants, $\mathbf{X}_n \xrightarrow{d} \mathbf{a} \Rightarrow \mathbf{X}_n \xrightarrow{P} \mathbf{a}$.
- Strong Law of Large Numbers (SLLN): Let $\mathbf{X}_1, \dots, \mathbf{X}_n$ be i.i.d. random vectors with finite first moment. Then $\bar{\mathbf{X}}_n \xrightarrow{a.s.} E(\mathbf{X}_1)$.
- Central Limit Theorem: Let $\mathbf{X}_1, \dots, \mathbf{X}_n$ be i.i.d. random vectors with expected value vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$. Then $\sqrt{n}(\bar{\mathbf{X}}_n - \boldsymbol{\mu})$ converges in distribution to a multivariate normal with mean $\mathbf{0}$ and covariance matrix $\boldsymbol{\Sigma}$.
- Slutsky Theorems for Convergence in Distribution:
 1. If $\mathbf{X}_n \in \mathbb{R}^m$, $\mathbf{X}_n \xrightarrow{d} \mathbf{X}$ and if $f : \mathbb{R}^m \rightarrow \mathbb{R}^q$ (where $q \leq m$) is continuous except possibly on a set C with $P(\mathbf{X} \in C) = 0$, then $f(\mathbf{X}_n) \xrightarrow{d} f(\mathbf{X})$.
 2. If $\mathbf{X}_n \xrightarrow{d} \mathbf{X}$ and $(\mathbf{X}_n - \mathbf{Y}_n) \xrightarrow{P} \mathbf{0}$, then $\mathbf{Y}_n \xrightarrow{d} \mathbf{X}$.
 3. If $\mathbf{X}_n \in \mathbb{R}^d$, $\mathbf{Y}_n \in \mathbb{R}^k$, $\mathbf{X}_n \xrightarrow{d} \mathbf{X}$ and $\mathbf{Y}_n \xrightarrow{d} \mathbf{c}$, then

$$\begin{pmatrix} \mathbf{X}_n \\ \mathbf{Y}_n \end{pmatrix} \xrightarrow{d} \begin{pmatrix} \mathbf{X} \\ \mathbf{c} \end{pmatrix}$$
- Slutsky Theorems for Convergence in Probability:
 1. If $\mathbf{X}_n \in \mathbb{R}^m$, $\mathbf{X}_n \xrightarrow{P} \mathbf{X}$ and if $f : \mathbb{R}^m \rightarrow \mathbb{R}^q$ (where $q \leq m$) is continuous except possibly on a set C with $P(\mathbf{X} \in C) = 0$, then $f(\mathbf{X}_n) \xrightarrow{P} f(\mathbf{X})$.
 2. If $\mathbf{X}_n \xrightarrow{P} \mathbf{X}$ and $(\mathbf{X}_n - \mathbf{Y}_n) \xrightarrow{P} \mathbf{0}$, then $\mathbf{Y}_n \xrightarrow{P} \mathbf{X}$.
 3. If $\mathbf{X}_n \in \mathbb{R}^d$, $\mathbf{Y}_n \in \mathbb{R}^k$, $\mathbf{X}_n \xrightarrow{P} \mathbf{X}$ and $\mathbf{Y}_n \xrightarrow{P} \mathbf{Y}$, then

$$\begin{pmatrix} \mathbf{X}_n \\ \mathbf{Y}_n \end{pmatrix} \xrightarrow{P} \begin{pmatrix} \mathbf{X} \\ \mathbf{Y} \end{pmatrix}$$
- Delta Method (Theorem of Cramér, Ferguson p. 45): Let $g : \mathbb{R}^d \rightarrow \mathbb{R}^k$ be such that the elements of $\dot{g}(\mathbf{x}) = \left[\frac{\partial g_i}{\partial x_j} \right]_{k \times d}$ are continuous in a neighborhood of $\boldsymbol{\theta} \in \mathbb{R}^d$. If \mathbf{X}_n is a sequence of d -dimensional random vectors such that $\sqrt{n}(\mathbf{X}_n - \boldsymbol{\theta}) \xrightarrow{d} \mathbf{X}$, then $\sqrt{n}(g(\mathbf{X}_n) - g(\boldsymbol{\theta})) \xrightarrow{d} \dot{g}(\boldsymbol{\theta})\mathbf{X}$. In particular, if $\sqrt{n}(\mathbf{X}_n - \boldsymbol{\theta}) \xrightarrow{d} \mathbf{X} \sim N(\mathbf{0}, \boldsymbol{\Sigma})$, then $\sqrt{n}(g(\mathbf{X}_n) - g(\boldsymbol{\theta})) \xrightarrow{d} \mathbf{Y} \sim N(\mathbf{0}, \dot{g}(\boldsymbol{\theta})\boldsymbol{\Sigma}\dot{g}(\boldsymbol{\theta})')$.

¹This material can be found in many texts. I took it from T. S. Ferguson's *A course in large sample theory*: Chapman and Hall, 1996