

The Multivariate Normal Distribution¹

STA 302 Fall 2016

¹See last slide for copyright information.

Overview

- 1 Moment-generating Functions
- 2 Definition
- 3 Properties
- 4 χ^2 and t distributions

Joint moment-generating function

Of a p -dimensional random vector \mathbf{x}

- $M_{\mathbf{x}}(\mathbf{t}) = E\left(e^{\mathbf{t}'\mathbf{x}}\right)$
- For example, $M_{(x_1, x_2, x_3)}(t_1, t_2, t_3) = E\left(e^{x_1 t_1 + x_2 t_2 + x_3 t_3}\right)$

Section 4.3 of *Linear models in statistics* has some material on moment-generating functions (optional).

Two big theorems

Proof omitted

- 1 Joint moment-generating functions correspond uniquely to joint probability distributions.
- 2 Two random vectors \mathbf{x}_1 and \mathbf{x}_2 are independent if and only if the moment-generating function of their joint distribution is the product of their moment-generating functions.

These results assume only that the moment-generating functions exist in a neighborhood of $\mathbf{t} = \mathbf{0}$. Nothing else is required.

A helpful distinction

- If x_1 and x_2 are independent,

$$M_{x_1+x_2}(t) = M_{x_1}(t)M_{x_2}(t)$$

- x_1 and x_2 are independent if and only if

$$M_{x_1, x_2}(t_1, t_2) = M_{x_1}(t_1)M_{x_2}(t_2)$$

Theorem: Functions of independent random vectors are independent

Show \mathbf{x}_1 and \mathbf{x}_2 independent implies that $\mathbf{y}_1 = g_1(\mathbf{x}_1)$ and $\mathbf{y}_2 = g_2(\mathbf{x}_2)$ are independent.

Let

$$\mathbf{y} = \begin{pmatrix} \mathbf{y}_1 \\ \mathbf{y}_2 \end{pmatrix} = \begin{pmatrix} g_1(\mathbf{x}_1) \\ g_2(\mathbf{x}_2) \end{pmatrix} \text{ and } \mathbf{t} = \begin{pmatrix} \mathbf{t}_1 \\ \mathbf{t}_2 \end{pmatrix}. \text{ Then}$$

$$\begin{aligned} M_{\mathbf{y}}(\mathbf{t}) &= E\left(e^{\mathbf{t}'\mathbf{y}}\right) \\ &= E\left(e^{\mathbf{t}'_1\mathbf{y}_1 + \mathbf{t}'_2\mathbf{y}_2}\right) = E\left(e^{\mathbf{t}'_1\mathbf{y}_1} e^{\mathbf{t}'_2\mathbf{y}_2}\right) \\ &= E\left(e^{\mathbf{t}'_1 g_1(\mathbf{x}_1)} e^{\mathbf{t}'_2 g_2(\mathbf{x}_2)}\right) \\ &= \int \int e^{\mathbf{t}'_1 g_1(\mathbf{x}_1)} e^{\mathbf{t}'_2 g_2(\mathbf{x}_2)} f_{\mathbf{x}_1}(\mathbf{x}_1) f_{\mathbf{x}_2}(\mathbf{x}_2) d(\mathbf{x}_1) d(\mathbf{x}_2) \\ &= M_{g_1(\mathbf{x}_1)}(\mathbf{t}_1) M_{g_2(\mathbf{x}_2)}(\mathbf{t}_2) \\ &= M_{\mathbf{y}_1}(\mathbf{t}_1) M_{\mathbf{y}_2}(\mathbf{t}_2) \end{aligned}$$

So \mathbf{y}_1 and \mathbf{y}_2 are independent.

$$M_{A\mathbf{x}}(\mathbf{t}) = M_{\mathbf{x}}(A'\mathbf{t})$$

Analogue of $M_{ax}(t) = M_x(at)$

$$\begin{aligned} M_{A\mathbf{x}}(\mathbf{t}) &= E\left(e^{\mathbf{t}'A\mathbf{x}}\right) \\ &= E\left(e^{(A'\mathbf{t})'\mathbf{x}}\right) \\ &= M_{\mathbf{x}}(A'\mathbf{t}) \end{aligned}$$

Note that \mathbf{t} is the same length as $\mathbf{y} = A\mathbf{x}$: The number of rows in A .

$$M_{\mathbf{x}+\mathbf{c}}(\mathbf{t}) = e^{\mathbf{t}'\mathbf{c}}M_{\mathbf{x}}(\mathbf{t})$$

Analogue of $M_{x+c}(t) = e^{ct}M_x(t)$

$$\begin{aligned}M_{\mathbf{x}+\mathbf{c}}(\mathbf{t}) &= E\left(e^{\mathbf{t}'(\mathbf{x}+\mathbf{c})}\right) \\&= E\left(e^{\mathbf{t}'\mathbf{x}+\mathbf{t}'\mathbf{c}}\right) \\&= e^{\mathbf{t}'\mathbf{c}}E\left(e^{\mathbf{t}'\mathbf{x}}\right) \\&= e^{\mathbf{t}'\mathbf{c}}M_{\mathbf{x}}(\mathbf{t})\end{aligned}$$

Distributions may be defined in terms of moment-generating functions

Build up the multivariate normal from univariate normals.

- If $y \sim N(\mu, \sigma^2)$, then $M_y(t) = e^{\mu t + \frac{1}{2}\sigma^2 t^2}$
- Moment-generating functions correspond uniquely to probability distributions.
- So *define* a normal random variable with expected value μ and variance σ^2 as a random variable with moment-generating function $e^{\mu t + \frac{1}{2}\sigma^2 t^2}$.
- This has one surprising consequence ...

Degenerate random variables

A *degenerate* random variable has all the probability concentrated at a single value, say $Pr\{y = y_0\} = 1$. Then

$$\begin{aligned}M_y(t) &= E(e^{yt}) \\&= \sum_y e^{yt} p(y) \\&= e^{y_0 t} \cdot p(y_0) \\&= e^{y_0 t} \cdot 1 \\&= e^{y_0 t}\end{aligned}$$

If $Pr\{y = y_0\} = 1$, then $M_y(t) = e^{y_0 t}$

- This is of the form $e^{\mu t + \frac{1}{2}\sigma^2 t^2}$ with $\mu = y_0$ and $\sigma^2 = 0$.
- So $y \sim N(y_0, 0)$.
- That is, degenerate random variables are “normal” with variance zero.
- Call them *singular* normals.
- This will be surprisingly handy later.

Independent standard normals

Let $z_1, \dots, z_p \stackrel{i.i.d.}{\sim} N(0, 1)$.

$$\mathbf{z} = \begin{pmatrix} z_1 \\ \vdots \\ z_p \end{pmatrix}$$

$$E(\mathbf{z}) = \mathbf{0} \qquad cov(\mathbf{z}) = I_p$$

Moment-generating function of \mathbf{z}

Using $e^{\mu t + \frac{1}{2}\sigma^2 t^2}$

$$\begin{aligned}M_{\mathbf{z}}(\mathbf{t}) &= \prod_{j=1}^p M_{z_j}(t_j) \\ &= \prod_{j=1}^p e^{\frac{1}{2}t_j^2} \\ &= e^{\frac{1}{2}\sum_{j=1}^p t_j^2} \\ &= e^{\frac{1}{2}\mathbf{t}'\mathbf{t}}\end{aligned}$$

Transform \mathbf{z} to get a general multivariate normal

Remember: A non-negative definite means $\mathbf{v}'\mathbf{A}\mathbf{v} \geq 0$

Let Σ be a $p \times p$ symmetric *non-negative definite* matrix and $\boldsymbol{\mu} \in \mathbb{R}^p$. Let $\mathbf{y} = \Sigma^{1/2}\mathbf{z} + \boldsymbol{\mu}$.

- The elements of \mathbf{y} are linear combinations of independent standard normals.
- Linear combinations of normals should be normal.
- \mathbf{y} has a multivariate distribution.
- We'd like to call \mathbf{y} a *multivariate normal*.

Moment-generating function of $\mathbf{y} = \Sigma^{1/2}\mathbf{z} + \boldsymbol{\mu}$

Remember: $M_{A\mathbf{x}}(\mathbf{t}) = M_{\mathbf{x}}(A'\mathbf{t})$ and $M_{\mathbf{x}+\mathbf{c}}(\mathbf{t}) = e^{\mathbf{t}'\mathbf{c}}M_{\mathbf{x}}(\mathbf{t})$ and $M_{\mathbf{z}}(\mathbf{t}) = e^{\frac{1}{2}\mathbf{t}'\mathbf{t}}$

$$\begin{aligned}
 M_{\mathbf{y}}(\mathbf{t}) &= M_{\Sigma^{1/2}\mathbf{z}+\boldsymbol{\mu}}(\mathbf{t}) \\
 &= e^{\mathbf{t}'\boldsymbol{\mu}} M_{\Sigma^{1/2}\mathbf{z}}(\mathbf{t}) \\
 &= e^{\mathbf{t}'\boldsymbol{\mu}} M_{\mathbf{z}}(\Sigma^{1/2}'\mathbf{t}) \\
 &= e^{\mathbf{t}'\boldsymbol{\mu}} M_{\mathbf{z}}(\Sigma^{1/2}\mathbf{t}) \\
 &= e^{\mathbf{t}'\boldsymbol{\mu}} e^{\frac{1}{2}(\Sigma^{1/2}\mathbf{t})'(\Sigma^{1/2}\mathbf{t})} \\
 &= e^{\mathbf{t}'\boldsymbol{\mu}} e^{\frac{1}{2}\mathbf{t}'\Sigma^{1/2}\Sigma^{1/2}\mathbf{t}} \\
 &= e^{\mathbf{t}'\boldsymbol{\mu}} e^{\frac{1}{2}\mathbf{t}'\Sigma\mathbf{t}} \\
 &= e^{\mathbf{t}'\boldsymbol{\mu}+\frac{1}{2}\mathbf{t}'\Sigma\mathbf{t}}
 \end{aligned}$$

So *define* a multivariate normal random variable \mathbf{y} as one with moment-generating function $M_{\mathbf{y}}(\mathbf{t}) = e^{\mathbf{t}'\boldsymbol{\mu}} e^{\frac{1}{2}\mathbf{t}'\Sigma\mathbf{t}}$.

Compare univariate and multivariate normal moment-generating functions

Univariate $M_y(t) = e^{\mu t + \frac{1}{2}\sigma^2 t^2}$

Multivariate $M_{\mathbf{y}}(\mathbf{t}) = e^{\mathbf{t}'\boldsymbol{\mu}} e^{\frac{1}{2}\mathbf{t}'\boldsymbol{\Sigma}\mathbf{t}}$

So the univariate normal is a special case of the multivariate normal with $p = 1$.

Mean and covariance matrix

For a univariate normal, $E(y) = \mu$ and $Var(y) = \sigma^2$

Recall $\mathbf{y} = \Sigma^{1/2}\mathbf{z} + \boldsymbol{\mu}$.

$$\begin{aligned} E(\mathbf{y}) &= \boldsymbol{\mu} \\ cov(\mathbf{y}) &= \Sigma^{1/2} cov(\mathbf{z}) \Sigma^{1/2'} \\ &= \Sigma^{1/2} I \Sigma^{1/2} \\ &= \Sigma \end{aligned}$$

We will say \mathbf{y} is multivariate normal with expected value $\boldsymbol{\mu}$ and variance-covariance matrix Σ , and write $\mathbf{y} \sim N_p(\boldsymbol{\mu}, \Sigma)$.

Note that because $M_{\mathbf{y}}(\mathbf{t}) = e^{\mathbf{t}'\boldsymbol{\mu}} e^{\frac{1}{2}\mathbf{t}'\Sigma\mathbf{t}}$, $\boldsymbol{\mu}$ and Σ completely determine the distribution.

Probability density function of $\mathbf{y} \sim N_p(\boldsymbol{\mu}, \Sigma)$

Remember, Σ is only positive *semi*-definite.

It is easy to write down the density of $\mathbf{z} \sim N_p(\mathbf{0}, I)$ as a product of standard normals.

If Σ is strictly positive definite (and not otherwise), the density of $\mathbf{y} = \Sigma^{1/2}\mathbf{z} + \boldsymbol{\mu}$ can be obtained using the Jacobian Theorem as

$$f(\mathbf{y}) = \frac{1}{|\Sigma|^{1/2} (2\pi)^{p/2}} \exp \left\{ -\frac{1}{2} (\mathbf{y} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{y} - \boldsymbol{\mu}) \right\}$$

This is usually how the multivariate normal is defined.

Σ positive definite?

- Positive definite means that for any non-zero $p \times 1$ vector \mathbf{a} , we have $\mathbf{a}'\Sigma\mathbf{a} > 0$.
- Since the one-dimensional random variable $w = \sum_{i=1}^p a_i y_i$ may be written as $W = \mathbf{a}'\mathbf{y}$ and $Var(w) = cov(\mathbf{a}'\mathbf{y}) = \mathbf{a}'\Sigma\mathbf{a}$, it is natural to require that Σ be positive definite.
- All it means is that every non-zero linear combination of \mathbf{y} values has a positive variance. Often, this is what you want.

Singular normal: Σ is positive *semi*-definite.

Suppose there is $\mathbf{a} \neq \mathbf{0}$ with $\mathbf{a}'\Sigma\mathbf{a} = 0$. Let $w = \mathbf{a}'\mathbf{y}$.

- Then $Var(w) = Var(\mathbf{a}'\mathbf{y}) = \mathbf{a}'\Sigma\mathbf{a} = 0$. That is, w has a degenerate distribution (but it's still normal).
- In this case we describe the distribution of \mathbf{y} as a *singular* multivariate normal.
- Including the singular case saves a lot of extra work in later proofs.
- We will insist that a singular multivariate normal is still multivariate normal, even though it has no density.

Distribution of $A\mathbf{y}$

Recall $\mathbf{y} \sim N_p(\boldsymbol{\mu}, \Sigma)$ means $M_{\mathbf{y}}(\mathbf{t}) = e^{\mathbf{t}'\boldsymbol{\mu} + \frac{1}{2}\mathbf{t}'\Sigma\mathbf{t}}$

Let $\mathbf{y} \sim N_p(\boldsymbol{\mu}, \Sigma)$, and $\mathbf{w} = A\mathbf{y}$, where A is an $r \times p$ matrix.

$$\begin{aligned}M_{\mathbf{w}}(\mathbf{t}) &= M_{A\mathbf{y}}(\mathbf{t}) \\&= M_{\mathbf{y}}(A'\mathbf{t}) \\&= e^{(A'\mathbf{t})'\boldsymbol{\mu}} e^{\frac{1}{2}(A'\mathbf{t})'\Sigma(A'\mathbf{t})} \\&= e^{\mathbf{t}'(A\boldsymbol{\mu})} e^{\frac{1}{2}\mathbf{t}'(A\Sigma A')\mathbf{t}} \\&= e^{\mathbf{t}'(A\boldsymbol{\mu}) + \frac{1}{2}\mathbf{t}'(A\Sigma A')\mathbf{t}}\end{aligned}$$

Recognize moment-generating function and conclude

$$\mathbf{w} \sim N_r(A\boldsymbol{\mu}, A\Sigma A')$$

Exercise

Use moment-generating functions, of course.

Let $\mathbf{y} \sim N_p(\boldsymbol{\mu}, \Sigma)$.

Show $\mathbf{y} + \mathbf{c} \sim N_p(\boldsymbol{\mu} + \mathbf{c}, \Sigma)$.

Zero covariance implies independence for the multivariate normal.

- Independence always implies zero covariance.
- For the multivariate normal, zero covariance also implies independence.
- The multivariate normal is the only continuous distribution with this property.

Show zero covariance implies independence

By showing $M_{\mathbf{y}}(\mathbf{t}) = M_{y_1}(\mathbf{t}_1)M_{y_2}(\mathbf{t}_2)$

Let $\mathbf{y} \sim N(\boldsymbol{\mu}, \Sigma)$, with

$$\mathbf{y} = \begin{pmatrix} \mathbf{y}_1 \\ \mathbf{y}_2 \end{pmatrix} \quad \boldsymbol{\mu} = \begin{pmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{pmatrix} \quad \Sigma = \begin{pmatrix} \Sigma_1 & \mathbf{0} \\ \mathbf{0} & \Sigma_2 \end{pmatrix} \quad \mathbf{t} = \begin{pmatrix} \mathbf{t}_1 \\ \mathbf{t}_2 \end{pmatrix}$$

$$\begin{aligned} M_{\mathbf{y}}(\mathbf{t}) &= E\left(e^{\mathbf{t}'\mathbf{y}}\right) \\ &= E\left(e^{\left(\frac{\mathbf{t}_1}{\mathbf{t}_2}\right)' \mathbf{y}}\right) \\ &= \dots \end{aligned}$$

Continuing the calculation: $M_{\mathbf{y}}(\mathbf{t}) = e^{\mathbf{t}'\boldsymbol{\mu} + \frac{1}{2}\mathbf{t}'\boldsymbol{\Sigma}\mathbf{t}}$

$$\mathbf{y} = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \quad \boldsymbol{\mu} = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix} \quad \boldsymbol{\Sigma} = \begin{pmatrix} \Sigma_1 & \mathbf{0} \\ \mathbf{0} & \Sigma_2 \end{pmatrix} \quad \mathbf{t} = \begin{pmatrix} t_1 \\ t_2 \end{pmatrix}$$

$$\begin{aligned} M_{\mathbf{y}}(\mathbf{t}) &= E \left(e^{\begin{pmatrix} t_1 \\ t_2 \end{pmatrix}' \mathbf{y}} \right) \\ &= \exp \left\{ (t_1' | t_2') \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix} \right\} \exp \left\{ \frac{1}{2} (t_1' | t_2') \begin{pmatrix} \Sigma_1 & \mathbf{0} \\ \mathbf{0} & \Sigma_2 \end{pmatrix} \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} \right\} \\ &= e^{t_1'\mu_1 + t_2'\mu_2} \exp \left\{ \frac{1}{2} (t_1'\Sigma_1 | t_2'\Sigma_2) \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} \right\} \\ &= e^{t_1'\mu_1 + t_2'\mu_2} \exp \left\{ \frac{1}{2} (t_1'\Sigma_1 t_1 + t_2'\Sigma_2 t_2) \right\} \\ &= e^{t_1'\mu_1} e^{t_2'\mu_2} e^{\frac{1}{2}(t_1'\Sigma_1 t_1)} e^{\frac{1}{2}(t_2'\Sigma_2 t_2)} \\ &= e^{t_1'\mu_1 + \frac{1}{2}(t_1'\Sigma_1 t_1)} e^{t_2'\mu_2 + \frac{1}{2}(t_2'\Sigma_2 t_2)} \\ &= M_{\mathbf{y}_1}(t_1) M_{\mathbf{y}_2}(t_2) \end{aligned}$$

So \mathbf{y}_1 and \mathbf{y}_2 are independent.

An easy example

If you do it the easy way

Let $y_1 \sim N(1, 2)$, $y_2 \sim N(2, 4)$ and $y_3 \sim N(6, 3)$ be independent, with $w_1 = y_1 + y_2$ and $w_2 = y_2 + y_3$. Find the joint distribution of w_1 and w_2 .

$$\begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

$$\mathbf{w} = A\mathbf{y} \sim N(A\boldsymbol{\mu}, A\Sigma A')$$

$$\mathbf{w} = A\mathbf{y} \sim N(A\boldsymbol{\mu}, A\Sigma A')$$

$y_1 \sim N(1, 2)$, $y_2 \sim N(2, 4)$ and $y_3 \sim N(6, 3)$ are independent

$$A\boldsymbol{\mu} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} = \begin{pmatrix} 3 \\ 8 \end{pmatrix}$$

$$\begin{aligned} A\Sigma A' &= \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 6 & 4 \\ 4 & 7 \end{pmatrix} \end{aligned}$$

Marginal distributions are multivariate normal

 $\mathbf{y} \sim N_p(\boldsymbol{\mu}, \Sigma)$, so $\mathbf{w} = A\mathbf{y} \sim N(A\boldsymbol{\mu}, A\Sigma A')$

Find the distribution of

$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{pmatrix} = \begin{pmatrix} y_2 \\ y_4 \end{pmatrix}$$

Bivariate normal. The expected value is easy.

Covariance matrix

$$\begin{aligned}
\text{cov}(\mathbf{A}\mathbf{y}) &= \mathbf{A}\Sigma\mathbf{A}' \\
&= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \sigma_1^2 & \sigma_{1,2} & \sigma_{1,3} & \sigma_{1,4} \\ \sigma_{1,2} & \sigma_2^2 & \sigma_{2,3} & \sigma_{2,4} \\ \sigma_{1,3} & \sigma_{2,3} & \sigma_3^2 & \sigma_{3,4} \\ \sigma_{1,4} & \sigma_{2,4} & \sigma_{3,4} & \sigma_4^2 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix} \\
&= \begin{pmatrix} \sigma_{1,2} & \sigma_2^2 & \sigma_{2,3} & \sigma_{2,4} \\ \sigma_{1,4} & \sigma_{2,4} & \sigma_{3,4} & \sigma_4^2 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix} \\
&= \begin{pmatrix} \sigma_2^2 & \sigma_{2,4} \\ \sigma_{2,4} & \sigma_4^2 \end{pmatrix}
\end{aligned}$$

Marginal distributions of a multivariate normal are multivariate normal, with the original means, variances and covariances.

Summary

- If \mathbf{c} is a vector of constants, $\mathbf{x} + \mathbf{c} \sim N(\mathbf{c} + \boldsymbol{\mu}, \Sigma)$
- If A is a matrix of constants, $A\mathbf{x} \sim N(A\boldsymbol{\mu}, A\Sigma A')$
- Linear combinations of multivariate normals are multivariate normal.
- All the marginals (dimension less than p) of \mathbf{x} are (multivariate) normal, but it is possible in theory to have a collection of univariate normals whose joint distribution is not multivariate normal.
- For the multivariate normal, zero covariance implies independence. The multivariate normal is the only continuous distribution with this property.

Showing $(\mathbf{x} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \sim \chi^2(p)$

$\boldsymbol{\Sigma}$ has to be positive definite this time

$$\begin{aligned}\mathbf{x} &\sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \\ \mathbf{y} = \mathbf{x} - \boldsymbol{\mu} &\sim N(\mathbf{0}, \boldsymbol{\Sigma}) \\ \mathbf{z} = \boldsymbol{\Sigma}^{-\frac{1}{2}} \mathbf{y} &\sim N\left(\mathbf{0}, \boldsymbol{\Sigma}^{-\frac{1}{2}} \boldsymbol{\Sigma} \boldsymbol{\Sigma}^{-\frac{1}{2}}\right) \\ &= N\left(\mathbf{0}, \boldsymbol{\Sigma}^{-\frac{1}{2}} \boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\Sigma}^{\frac{1}{2}} \boldsymbol{\Sigma}^{-\frac{1}{2}}\right) \\ &= N(\mathbf{0}, I)\end{aligned}$$

So \mathbf{z} is a vector of p independent standard normals, and

$$\mathbf{y}' \boldsymbol{\Sigma}^{-1} \mathbf{y} = (\boldsymbol{\Sigma}^{-\frac{1}{2}} \mathbf{y})' (\boldsymbol{\Sigma}^{-\frac{1}{2}} \mathbf{y}) = \mathbf{z}' \mathbf{z} = \sum_{j=1}^p z_j^2 \sim \chi^2(p) \quad \blacksquare$$

\bar{x} and s^2 independent

$$x_1, \dots, x_n \stackrel{i.i.d.}{\sim} N(\mu, \sigma^2)$$

$$\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \sim N(\mu \mathbf{1}, \sigma^2 I) \quad \mathbf{y} = \begin{pmatrix} x_1 - \bar{x} \\ \vdots \\ x_n - \bar{x} \\ \bar{x} \end{pmatrix} = A\mathbf{x}$$

Note A is $(n+1) \times n$, so $\text{cov}(A\mathbf{x}) = \sigma^2 \mathbf{A}\mathbf{A}'$ is $(n+1) \times (n+1)$, singular.

The argument

$$\mathbf{y} = A\mathbf{x} = \begin{pmatrix} x_1 - \bar{x} \\ \vdots \\ x_n - \bar{x} \\ \bar{x} \end{pmatrix} = \begin{pmatrix} \mathbf{y}_2 \\ \hline \bar{x} \end{pmatrix}$$

- \mathbf{y} is multivariate normal.
- $Cov(\bar{x}, (x_j - \bar{x})) = 0$ (Exercise)
- So \bar{x} and \mathbf{y}_2 are independent.
- So \bar{x} and $S^2 = g(\mathbf{y}_2)$ are independent. ■

Leads to the t distribution

If

- $z \sim N(0, 1)$ and
- $y \sim \chi^2(\nu)$ and
- z and y are independent, then

$$T = \frac{z}{\sqrt{y/\nu}} \sim t(\nu)$$

Random sample from a normal distribution

Let $x_1, \dots, x_n \stackrel{i.i.d.}{\sim} N(\mu, \sigma^2)$. Then

- $\frac{\sqrt{n}(\bar{x} - \mu)}{\sigma} \sim N(0, 1)$ and
- $\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$ and
- These quantities are independent, so

$$\begin{aligned} T &= \frac{\sqrt{n}(\bar{x} - \mu)/\sigma}{\sqrt{\frac{(n-1)S^2}{\sigma^2}/(n-1)}} \\ &= \frac{\sqrt{n}(\bar{x} - \mu)}{S} \sim t(n-1) \end{aligned}$$

Copyright Information

This slide show was prepared by **Jerry Brunner**, Department of Statistical Sciences, University of Toronto. It is licensed under a **Creative Commons Attribution - ShareAlike 3.0 Unported License**. Use any part of it as you like and share the result freely. The L^AT_EX source code is available from the course website:

<http://www.utstat.toronto.edu/~brunner/oldclass/302f16>