

STA 413F2011 Assignment 6

Do this assignment in preparation for Quiz Six in tutorial on Friday Oct. 21st. The problems are practice for the quiz, and are not to be handed in. See [Formula Sheet 4](#); any necessary material from the formula sheet will be supplied with the quiz.

Please take a look at Section 4.3.2. We proved the delta method another way in lecture. Also, please read pages 224-225. The following questions are practice for Test 3 and the Final Examination. They are not to be handed in.

1. Let $T_n \xrightarrow{P} \theta$, and let Y_n that is between T_n and θ . Prove that $Y_n \xrightarrow{P} \theta$. We will call this the Squeeze Theorem for Convergence in Probability.
2. The various parts of this question will lead you through the proof of the Delta Method for the special but important case of a function of the sample mean. That is, you are proving the result described as “Delta method combined with CLT” on the formula sheet. Of course, you may use anything on the formula sheet *except* the delta method.

(a) First, use Taylor’s Theorem (on the formula sheet) to re-write

$$\sqrt{n}(g(\bar{X}_n) - g(\mu)), \tag{1}$$

distributing \sqrt{n} . You now have two terms.

- (b) How do you know $\sqrt{n}(\bar{X}_n - \mu)$ converges in probability to a normal random variable? Cite something from the formula sheet.
 - (c) How do you know $g''(\mu^*) \xrightarrow{P} g''(\mu)$? Use the formula sheet as well as another problem from this assignment.
 - (d) Now show that the second of the two terms from Problem [2a](#) converges in probability to zero. When you use something from the formula sheet, cite it.
 - (e) Now show that the first term from Problem [2a](#) converges in distribution to something. What is its target distribution? Again, when you use something from the formula sheet, cite it.
 - (f) Finally, apply a rule from the formula sheet to establish the desired result. What is the rule? What is A_n ? What is B_n ?
3. Do Exercises 4.4.11 and 4.4.12.
 4. Let X_1, \dots, X_n be a random sample from a Bernoulli distribution with parameter θ . Find the limiting distribution of

$$Z_n = 2\sqrt{n} \left(\sin^{-1} \sqrt{\bar{X}_n} - \sin^{-1} \sqrt{\theta} \right).$$

Hint: $\frac{d}{dx} \sin^{-1}(x) = \frac{1}{\sqrt{1-x^2}}$.

5. Let X_1, \dots, X_n be a random sample from an exponential distribution with parameter θ .

- (a) Find a variance-stabilizing transformation. That is, find a function $g(x)$ such that the limiting distribution of

$$Y_n = \sqrt{n}[g(\bar{X}_n) - g(\theta)]$$

does not depend on θ .

- (b) To check, find the limiting distribution of Y_n .

6. Let X_1, \dots, X_n be a random sample from a uniform distribution on $(0, \theta)$.

- (a) Find a variance-stabilizing transformation. That is, find a function $g(x)$ such that the limiting distribution of

$$Y_n = \sqrt{n}[g(2\bar{X}_n) - g(\theta)]$$

does not depend on θ .

- (b) To check, find the limiting distribution of Y_n .

7. Let X_1, \dots, X_n be a random sample from a chi-square distribution with parameter ν .

- (a) Find a variance-stabilizing transformation. That is, find a function $g(x)$ such that the limiting distribution of

$$Y_n = \sqrt{n}[g(\bar{X}_n) - g(\nu)]$$

does not depend on ν .

- (b) To check, find the limiting distribution of Y_n .

8. Let X_1, \dots, X_n be a random sample from a binomial distribution with parameters m and θ . The constant m is fixed and known.

- (a) Find a variance-stabilizing transformation. That is, find a function $g(x)$ such that the limiting distribution of

$$Y_n = \sqrt{n}[g(\bar{X}_n/m) - g(\theta)]$$

does not depend on θ .

- (b) To check, find the limiting distribution of Y_n .