

# Chapter 5

## Multivariate and Within-cases Analysis

### 5.1 Multivariate Analysis of Variance

Multivariate means more than one dependent variable at once. Why do it? Primarily because if you do parallel analyses on lots of outcome measures, the probability of getting significant results just by chance will definitely exceed the apparent  $\alpha = 0.05$  level. It is also possible in principle to detect results from a multivariate analysis that are not significant at the univariate level.

The simplest way to do a multivariate analysis is to do a univariate analysis on each dependent variable separately, and apply a Bonferroni correction. The disadvantage is that testing this way is less powerful than doing it with real multivariate tests.

Another advantage of a true multivariate analysis is that it can “notice” things missed by several Bonferroni-corrected univariate analyses, because under the surface, a classical multivariate analysis involves the construction of the unique linear combination of the dependent variables that shows the strongest relationship (in the sense explaining the remaining variation) with the independent variables. The linear combination in question is called the first canonical variate or canonical variable.

- The number of canonical variables equals the number of dependent variables (or independent variables, whichever is fewer).
- The canonical variables are all uncorrelated with each other. The second one is constructed so that it has as strong a relationship as possible

to the independent variables – subject to the constraint that it have zero correlation with the first one, and so on.

- This why it is not optimal to do a principal components analysis (or factor analysis) on a set of dependent variables, and then treat the components (or factor scores) as dependent variables. Ordinary multivariate analysis is already doing this, and doing it much better.

### 5.1.1 Assumptions

As in the case of univariate analysis, the statistical assumptions of multivariate analysis concern conditional distributions – conditional upon various configurations of independent variable  $\mathbf{X}$  values. Here we are talking about the conditional joint distribution of several dependent variables observed for each case, say  $Y_1, \dots, Y_k$ . These are often described as a “vector” of observations. It may help to think of the collection of dependent variable values for a case as a point in  $k$ -dimensional space, and to imagine an arrow pointing from the origin  $(0, \dots, 0)$  to the point  $(Y_1, \dots, Y_k)$ ; the arrow is literally a vector. As I say, this may help. Or it may not.

The classical assumptions of multivariate analysis depend on the idea of a population covariance. The population covariance between  $Y_2$  and  $Y_4$  is denoted  $\sigma_{2,4}$ , and is defined by

$$\sigma_{2,4} = \rho_{2,4}\sigma_2\sigma_4,$$

where

$\sigma_2$  is the population standard deviation of  $Y_2$ ,

$\sigma_4$  is the population standard deviation of  $Y_4$ , and

$\rho_{2,4}$  is the population correlation between  $Y_2$  and  $Y_4$  (That’s the Greek letter rho).

The population covariance can be estimated by the sample covariance, defined in a parallel way by  $s_{2,4} = r_{2,4}s_2s_4$ , where  $s_2$  and  $s_4$  are the sample standard deviations and  $r$  is the Pearson correlation coefficient. Whether we are talking about population parameters or sample statistics, it is clear that zero covariance means zero correlation and vice versa.

We will use  $\Sigma$  (the capital Greek letter sigma) to stand for the population variance-covariance matrix. This is a  $k$  by  $k$  rectangular array of numbers with variances on the main diagonal, and covariances on the off-diagonals. For 4 dependent variables it would look like this:

$$\Sigma = \begin{bmatrix} \sigma_1^2 & \sigma_{1,2} & \sigma_{1,3} & \sigma_{1,4} \\ \sigma_{1,2} & \sigma_2^2 & \sigma_{2,3} & \sigma_{2,4} \\ \sigma_{1,3} & \sigma_{2,3} & \sigma_3^2 & \sigma_{3,4} \\ \sigma_{1,4} & \sigma_{2,4} & \sigma_{3,4} & \sigma_4^2 \end{bmatrix}.$$


Notice the symmetry: Element  $(i, j)$  of a covariance matrix equals element  $(j, i)$ .

With this background, the assumptions of classical multivariate analysis are parallel to those of the standard univariate analysis of variance. Conditionally on the independent variable values,

- Sample vectors  $\mathbf{Y} = (Y_1, \dots, Y_k)$  represent *independent observations* for different cases.
- Each conditional distribution is *multivariate normal*.
- Each conditional distribution has the *same population variance-covariance matrix*.

The multivariate normal distribution is a generalization of the one-dimensional normal. Instead of probabilities being areas under a curve they are now volumes under a surface. Here is a picture of the bivariate normal density (for  $k = 2$  dependent variables).

Figure 5.1: Bivariate Normal Density



### 5.1.2 Significance Testing

In univariate analysis, different standard methods for deriving tests (these are hidden from you) all point to Fisher's  $F$  test. In multivariate analysis there are four major test statistics, *Wilks' Lambda*, *Pillai's Trace*, the *Hotelling-Lawley Trace*, and *Roy's Greatest Root*.

When there is only one dependent variable, these are all equivalent to  $F$ . When there is more than one dependent variable they are all about equally "good" (in any reasonable sense), and conclusions from them generally agree – but not always. Sometimes one will designate a finding as significant and another will not. In this case you have borderline results and there is no conventional way out of the dilemma.

The four multivariate test statistics all have  $F$  approximations that are used by SAS and other stat packages to compute p-values. Tables are available in textbooks on multivariate analysis. For the first three tests (Wilks' Lambda, Pillai's Trace and the Hotelling-Lawley Trace), the  $F$  approximations are very good. For Roy's greatest root the  $F$  approximation is lousy.

This is a problem with the cheap method for getting  $p$ -values, not with the test itself. One can always use tables.

When a multivariate test is significant, many people then follow up with ordinary univariate tests to see "which dependent variable the results came from." This is a reasonable exploratory strategy. More conservative is to follow up with Bonferroni-corrected univariate tests. When you do this, however, there is no guarantee that any of the Bonferroni-corrected tests will be significant.

It is also possible, and in some ways very appealing, to follow up a significant multivariate test with Scheffé tests. For example, Scheffé follow-ups to a significant one-way multivariate ANOVA would include adjusted versions of all the corresponding univariate one-way ANOVAs, all multivariate pairwise comparisons, all univariate pairwise comparisons, and countless other possibilities — all simultaneously protected at the 0.05 level.

You can also try interpret a significant multivariate effect by looking at the canonical variates, but there is no guarantee they will make sense.

### 5.1.3 The Hospital Example

In the following example, cases are hospitals in 4 different regions of the U.S.. The hospitals either have a medical school affiliation or not. The dependent variables are average length of time a patient stays at the hospital, and infection risk – the estimated probability that a patient will acquire an infection unrelated to what he or she came in with. We will analyze these data as a two-way multivariate analysis of variance.

```
/****** senicmv96a.sas *****/
title 'Senic data: SAS glm & reg multivariate intro';

%include 'senicdef.sas'; /* senicdef.sas reads data, etc.
                          Includes reg1-reg3, ms1 & mr1-mr3 */

proc glm;
  class region medschl;
  model infrisk stay = region|medschl;
  manova h = _all_;
```

The `proc glm` output starts with full univariate output for each dependent variable. Then (for each effect tested) there is some multivariate output you ignore,

General Linear Models Procedure  
Multivariate Analysis of Variance

Characteristic Roots and Vectors of: E Inverse \* H, where  
H = Type III SS&CP Matrix for REGION    E = Error SS&CP Matrix

Characteristic Root	Percent	Characteristic Vector	V'EV=1
		INFRISK	STAY
0.14830859	95.46	-0.00263408	0.06067199
0.00705986	4.54	0.08806967	-0.03251114

followed by the interesting part.

Manova Test Criteria and F Approximations for  
the Hypothesis of no Overall REGION Effect  
H = Type III SS&CP Matrix for REGION    E = Error SS&CP Matrix

	S=2	M=0	N=51	Value	F	Num DF	Den DF	Pr > F
Statistic								
Wilks' Lambda				0.86474110	2.6127	6	208	0.0183
Pillai's Trace				0.13616432	2.5570	6	210	0.0207
Hotelling-Lawley Trace				0.15536845	2.6672	6	206	0.0163
Roy's Greatest Root				0.14830859	5.1908	3	105	0.0022

NOTE: F Statistic for Roy's Greatest Root is an upper bound.  
NOTE: F Statistic for Wilks' Lambda is exact.

...

Manova Test Criteria and Exact F Statistics for

the Hypothesis of no Overall MEDSCHL Effect  
H = Type III SS&CP Matrix for MEDSCHL E = Error SS&CP Matrix

	S=1	M=0	N=51			
Statistic	Value	F	Num DF	Den DF	Pr > F	
Wilks' Lambda	0.92228611	4.3816	2	104	0.0149	
Pillai's Trace	0.07771389	4.3816	2	104	0.0149	
Hotelling-Lawley Trace	0.08426224	4.3816	2	104	0.0149	
Roy's Greatest Root	0.08426224	4.3816	2	104	0.0149	

NOTE: F Statistic for Roy's Greatest Root is an upper bound.

. . . .

Manova Test Criteria and F Approximations for  
the Hypothesis of no Overall REGION\*MEDSCHL Effect  
H = Type III SS&CP Matrix for REGION\*MEDSCHL E = Error SS&CP Matrix

	S=2	M=0	N=51			
Statistic	Value	F	Num DF	Den DF	Pr > F	
Wilks' Lambda	0.95784589	0.7546	6	208	0.6064	
Pillai's Trace	0.04228179	0.7559	6	210	0.6054	
Hotelling-Lawley Trace	0.04387599	0.7532	6	206	0.6075	
Roy's Greatest Root	0.04059215	1.4207	3	105	0.2409	

NOTE: F Statistic for Roy's Greatest Root is an upper bound.

NOTE: F Statistic for Wilks' Lambda is exact.

Remember the output started with the univariate analyses. We'll look at them here (out of order) – just Type III SS, because that's parallel to the multivariate tests. We are tracking down the significant multivariate effects for Region and Medical School Affiliation. Using Bonferroni correction means only believe it if  $p < 0.025$ .

Dependent Variable: INFRISK prob of acquiring infection in hospital

Source	DF	Type III SS	Mean Square	F Value	Pr > F
REGION	3	6.61078342	2.20359447	1.35	0.2623
MEDSCHL	1	6.64999500	6.64999500	4.07	0.0461
REGION*MEDSCHL	3	5.32149160	1.77383053	1.09	0.3581

Dependent Variable: STAY av length of hospital stay, in days

Source	DF	Type III SS	Mean Square	F Value	Pr > F
REGION	3	41.61422755	13.87140918	5.19	0.0022
MEDSCHL	1	22.49593643	22.49593643	8.41	0.0045
REGION*MEDSCHL	3	0.92295998	0.30765333	0.12	0.9511

We conclude that the multivariate effect comes from a univariate relationship between the independent variables and stay. Question: If this is what we were going to do in the end, why do a multivariate analysis at all? Why not just two univariate analyses with a Bonferroni correction?

## 5.2 Within-cases (Repeated Measures) Analysis of Variance

In certain kinds of experimental research, it is common to obtain repeated measurements of a variable from the same individual at several different points in time. Usually it is unrealistic to assume that these repeated observations are uncorrelated, and it is very desirable to build their inter-correlations into the statistical model.

Sometimes, an individual (in some combination of experimental conditions) is measured under essentially the same conditions at several different points in time. In that case we will say that time is a *within-subjects* factor, because each subject contributes data at more than one value of the independent variable “time.” If a subject experiences only one value of an independent variable, it is called a between subjects factor.



Sometimes, an individual experiences more than one experimental treatment — for example judging the same stimuli under different background noise levels. In this case, the order of presentation of different noise levels would be counterbalanced so that time and noise level are unrelated (not confounded). Here noise level would be a within-subjects factor. The same study can definitely have more than one within-subjects factor and more than one between subjects factor.

The meaning of main effects and interactions, as well as their graphical presentation, is the same for within and between subjects factors.

We will discuss three methods for analyzing repeated measures data. In an order that is convenient but not historically chronological they are

1. The multivariate approach
2. The classical univariate approach
3. The covariance structure approach

### 5.2.1 The multivariate approach to repeated measures

First, note that any of the 3 methods can be multivariate, in the sense that several dependent variables can be measured at more than one time point. We will start with the simple case in which a single dependent variable is measured for each subject on several different occasions.

The basis of the multivariate approach to repeated measures is that *the different measurements conducted on each individual should be considered as multiple dependent variables*.

If there are  $k$  dependent variables, regular multivariate analysis allows for the analysis of up to  $k$  linear combinations of those dependent variables, instead of the original dependent variables. The multivariate approach to repeated measures sets up those linear combinations to be meaningful in terms of representing the within-cases structure of the data.

For example, suppose that men and women in 3 different age groups are tested on their ability to detect a signal under 5 different levels of background noise. There are 10 women and 10 men in each age group for a total  $n = 60$ . Order of presentation of noise levels is randomized for each subject, and the subjects themselves are tested in random order. This is a three-factor design. Age and sex are between subjects factors, and noise level is a within-subjects factor.

Let  $Y_1, Y_2, Y_3, Y_4$  and  $Y_5$  be the “Detection Scores” under the 5 different noise levels. Their population means are  $\mu_1, \mu_2, \mu_3, \mu_4$  and  $\mu_5$ , respectively.

We now construct 5 linear combinations of the Y variables, and give their

$$\begin{array}{ll}
 W_1 = (Y_1 + Y_2 + Y_3 + Y_4 + Y_5)/5 & E(W_1) = (\mu_1 + \mu_2 + \mu_3 + \mu_4 + \mu_5)/5 \\
 W_2 = Y_1 - Y_2 & E(W_2) = \mu_1 - \mu_2 \\
 W_3 = Y_2 - Y_3 & E(W_3) = \mu_2 - \mu_3 \\
 W_4 = Y_3 - Y_4 & E(W_4) = \mu_3 - \mu_4 \\
 W_5 = Y_4 - Y_5 & E(W_5) = \mu_4 - \mu_5
 \end{array}$$

expected values (population means).

Tests for main effects and interactions are obtained by treating these linear combinations (the  $W$ s) as dependent variables.

**Between-subjects effects** The main effects for age and sex, and the age by sex interaction, are just analyses conducted as usual on a single linear combination of the dependent variables, that is, on  $W_1$ . This is what we want; we are just averaging across within-subject values.

**Within-subject effects** Suppose that (averaging across treatment groups)  $E(W_2) = E(W_3) = E(W_4) = E(W_5) = 0$ . This means  $\mu_1 = \mu_2, \mu_2 = \mu_3, \mu_3 = \mu_4$  and  $\mu_4 = \mu_5$ . That is, there is no difference among noise level means, i.e., no main effect for the within-subjects factor.

**Interactions** of between and within-subjects factors are between-subjects effects tested simultaneously on the dependent variables representing differences among within-subject values –  $W_2$  through  $W_5$  in this case. For example, a significant sex difference in  $W_2$  through  $W_5$  means that the pattern of differences in mean discrimination among noise levels is different for males and females. Conceptually, this is exactly a noise level by sex interaction.

Similarly, a sex by age interaction on  $W_2$  through  $W_5$  means that the pattern of differences in mean discrimination among noise levels depends on special combinations of age and sex – a three-way (age by sex by noise) interaction.

## 5.2.2 The Noise Example

Here is noise.dat. Order of vars is ident, interest, sex, age, noise level, time noise level presented, discrim score. Notice that there are five lines of data for each case.

```

1 2.5 1 2 1 4 50.7
1 2.5 1 2 2 1 27.4
1 2.5 1 2 3 3 39.1
1 2.5 1 2 4 2 37.5
1 2.5 1 2 5 5 35.4
2 1.9 1 2 1 3 40.3
2 1.9 1 2 2 1 30.1
2 1.9 1 2 3 5 38.9
2 1.9 1 2 4 2 31.9
2 1.9 1 2 5 4 31.6
3 1.8 1 3 1 2 39.0
3 1.8 1 3 2 5 39.1
3 1.8 1 3 3 4 35.3
3 1.8 1 3 4 3 34.8
3 1.8 1 3 5 1 15.4
4 2.2 0 1 1 2 41.5
4 2.2 0 1 2 4 42.5

```

```

/***** noise96a.sas *****/
options linesize=79 pagesize=250;
title 'Repeated measures on Noise data: Multivariate approach';
proc format;      value sexfmt    0 = 'Male'  1 = 'Female' ;
data loud;
  infile 'noise.dat'; /* Multivariate data read */
  input ident  interest  sex  age  noise1 time1 discrim1
        ident2 inter2   sex2 age2 noise2 time2 discrim2
        ident3 inter3   sex3 age3 noise3 time3 discrim3
        ident4 inter4   sex4 age4 noise4 time4 discrim4
        ident5 inter5   sex5 age5 noise5 time5 discrim5 ;
  format sex sex2-sex5 sexfmt.;
  /* noise1 = 1, ... noise5 = 5. time1 = time noise 1 presented etc.
     ident, interest, sex & age are identical on each line */
  label interest = 'Interest in topic (politics)';

proc glm;
  class age sex;

```

```

model discrim1-discrim5 = age|sex;
repeated noise profile/ short summary;

```

First we get univariate analyses of discrim1-discrim5 – not the transformed vars yet. Then,

General Linear Models Procedure  
 Repeated Measures Analysis of Variance  
 Repeated Measures Level Information

Dependent Variable	DISCRIM1	DISCRIM2	DISCRIM3	DISCRIM4	DISCRIM5
Level of NOISE	1	2	3	4	5

Manova Test Criteria and Exact F Statistics for  
 the Hypothesis of no NOISE Effect  
 H = Type III SS&CP Matrix for NOISE    E = Error SS&CP Matrix

S=1    M=1    N=24.5

Statistic	Value	F	Num DF	Den DF	Pr > F
Wilks' Lambda	0.45363698	15.3562	4	51	0.0001
Pillai's Trace	0.54636302	15.3562	4	51	0.0001
Hotelling-Lawley Trace	1.20440581	15.3562	4	51	0.0001
Roy's Greatest Root	1.20440581	15.3562	4	51	0.0001

Manova Test Criteria and F Approximations for  
 the Hypothesis of no NOISE\*AGE Effect  
 H = Type III SS&CP Matrix for NOISE\*AGE    E = Error SS&CP Matrix

S=2    M=0.5    N=24.5

Statistic	Value	F	Num DF	Den DF	Pr > F
Wilks' Lambda	0.84653930	1.1076	8	102	0.3645
Pillai's Trace	0.15589959	1.0990	8	104	0.3700
Hotelling-Lawley Trace	0.17839904	1.1150	8	100	0.3597
Roy's Greatest Root	0.16044230	2.0857	4	52	0.0960

NOTE: F Statistic for Roy's Greatest Root is an upper bound.

NOTE: F Statistic for Wilks' Lambda is exact.

Manova Test Criteria and Exact F Statistics for  
the Hypothesis of no NOISE\*SEX Effect  
H = Type III SS&CP Matrix for NOISE\*SEX E = Error SS&CP Matrix

S=1 M=1 N=24.5

Statistic	Value	F	Num DF	Den DF	Pr > F
Wilks' Lambda	0.93816131	0.8404	4	51	0.5060
Pillai's Trace	0.06183869	0.8404	4	51	0.5060
Hotelling-Lawley Trace	0.06591477	0.8404	4	51	0.5060
Roy's Greatest Root	0.06591477	0.8404	4	51	0.5060

Manova Test Criteria and F Approximations for  
the Hypothesis of no NOISE\*AGE\*SEX Effect  
H = Type III SS&CP Matrix for NOISE\*AGE\*SEX E = Error SS&CP Matrix

S=2 M=0.5 N=24.5

Statistic	Value	F	Num DF	Den DF	Pr > F
Wilks' Lambda	0.84817732	1.0942	8	102	0.3735
Pillai's Trace	0.15679252	1.1058	8	104	0.3654
Hotelling-Lawley Trace	0.17313932	1.0821	8	100	0.3819
Roy's Greatest Root	0.12700316	1.6510	4	52	0.1755

NOTE: F Statistic for Roy's Greatest Root is an upper bound.

NOTE: F Statistic for Wilks' Lambda is exact.

General Linear Models Procedure  
Repeated Measures Analysis of Variance  
Tests of Hypotheses for Between Subjects Effects

Source	DF	Type III SS	Mean Square	F Value	Pr > F
AGE	2	1751.814067	875.907033	5.35	0.0076
SEX	1	77.419200	77.419200	0.47	0.4946
AGE*SEX	2	121.790600	60.895300	0.37	0.6911
Error	54	8839.288800	163.690533		

Then we are given "Univariate Tests of Hypotheses for Within Subject Effects" We will discuss these later. After that in the 1st file,

Repeated measures on Noise data: Multivariate approach

General Linear Models Procedure  
Repeated Measures Analysis of Variance  
Analysis of Variance of Contrast Variables

NOISE.N represents the nth successive difference in NOISE

Contrast Variable: NOISE.1

Source	DF	Type III SS	Mean Square	F Value	Pr > F
MEAN	1	537.00416667	537.00416667	5.40	0.0239
AGE	2	10.92133333	5.46066667	0.05	0.9466
SEX	1	45.93750000	45.93750000	0.46	0.4996
AGE*SEX	2	83.67600000	41.83800000	0.42	0.6587
Error	54	5370.09100000	99.44612963		

Contrast Variable: NOISE.2

Source	DF	Type III SS	Mean Square	F Value	Pr > F
MEAN	1	140.14816667	140.14816667	1.36	0.2489
AGE	2	106.89233333	53.44616667	0.52	0.5985
SEX	1	33.90016667	33.90016667	0.33	0.5688
AGE*SEX	2	159.32233333	79.66116667	0.77	0.4670
Error	54	5569.94700000	103.14716667		

Contrast Variable: NOISE.3

Source	DF	Type III SS	Mean Square	F Value	Pr > F
MEAN	1	50.41666667	50.41666667	0.72	0.4012
AGE	2	56.40633333	28.20316667	0.40	0.6720
SEX	1	195.84266667	195.84266667	2.78	0.1012
AGE*SEX	2	152.63633333	76.31816667	1.08	0.3456
Error	54	3802.61800000	70.41885185		

Contrast Variable: NOISE.4

Source	DF	Type III SS	Mean Square	F Value	Pr > F
MEAN	1	518.61600000	518.61600000	7.77	0.0073
AGE	2	449.45100000	224.72550000	3.37	0.0418
SEX	1	69.55266667	69.55266667	1.04	0.3118
AGE*SEX	2	190.97433333	95.48716667	1.43	0.2479
Error	54	3602.36600000	66.71048148		

### 5.2.3 The classical univariate approach to repeated measures

The univariate approach to repeated measures is chronologically the oldest. It can be derived in a clever way from the multivariate tests involving within subjects factors. It's what you get at the end of the default glm output – before the analysis of transformed variables, which you have to request specially.

General Linear Models Procedure  
Repeated Measures Analysis of Variance  
Univariate Tests of Hypotheses for Within Subject Effects

Source: NOISE

DF	Type III SS	Mean Square	F Value	Pr > F	Adj G - G	Pr > F H - F
4	2289.31400000	572.32850000	14.12	0.0001	0.0001	0.0001

Source: NOISE\*AGE

DF	Type III SS	Mean Square	F Value	Pr > F	Adj G - G	Pr > F H - F
8	334.42960000	41.80370000	1.03	0.4134	0.4121	0.4134

(The adj. G - G business will be explained later)

Source: NOISE\*SEX

DF	Type III SS	Mean Square	F Value	Pr > F	Adj G - G	Pr > F H - F
4	142.42280000	35.60570000	0.88	0.4777	0.4722	0.4777

Source: NOISE\*AGE\*SEX

DF	Type III SS	Mean Square	F Value	Pr > F	Adj G - G	Pr > F H - F
8	345.66440000	43.20805000	1.07	0.3882	0.3877	0.3882

Source: Error(NOISE)

DF	Type III SS	Mean Square
216	8755.83320000	40.53626481



Greenhouse-Geisser Epsilon = 0.9356  
Huynh-Feldt Epsilon = 1.1070

The classical univariate model for repeated measures is a mixed or sometimes a random effects model in which *subject* is a factor that, because its values are randomly sampled from a large population of potential subjects (just pretend), is a random factor. This factor is nested within any between-subjects factors; for example, Subject One in the “Male” group is a different person from Subject One in the “Female” group. The factor *subject* does not interact with any other factors. Interactions between subjects and various factors may sometimes be formally computed, but if they are computed they are always error terms; they are never tested.

In the noise level example, we could do

```
/****** noise96b.sas *****/
options linesize=79 pagesize=250;
title 'Repeated measures on Noise data: Univariate approach';
proc format;      value sexfmt    0 = 'Male'  1 = 'Female' ;

data loud;
  infile 'noise.dat'; /* Univariate data read */
  input ident interest sex age noise time discrim ;
  format sex sexfmt.;
  label interest = 'Interest in topic (politics)'
        time      = 'Order of presenting noise level';

proc glm;
  class age sex noise ident;
  model discrim = ident(age*sex) age|sex|noise;
  random ident(age*sex) / test;
```

- **Notice the univariate data read!** We are assuming  $n$  = number of observations, not number of cases.
- The results are identical to the univariate output produced as a by-product of the multivariate approach to repeated measures – if you know where to look.

- The overall (initial) test, and tests associated with Type I and Type III SS are all invalid.
- There are expected mean squares, which you should probably ignore.
- There are also repeated warnings that "This test assumes one or more other fixed effects are zero." SAS is buying testability of the hypotheses by assuming that you're only interested in an effect if all the higher-order interactions involving the effect are absent.

The univariate approach to repeated measures has some real virtues, sometimes. Because  $n$  = the number of observations rather than the number of cases, it is possible to have more parameters in a model than cases, or even more measurements than cases. In this situation the multivariate approach just blows up. It's either the classical univariate approach or the covariance structure approach, which will be discussed later.

The univariate approach may assume  $n$  is the number of observations, but it does not assume those observations are independent. In fact, the observations that come from the same subject are assumed to be correlated. The following discussion assumes that the model has no interactions between subjects and other factors; they're not only not tested, they're not even computed. This is not the only way to do it, and in fact sometimes the univariate tests produced by the `repeated` statement in `proc glm` are based on models with such interactions. But the strict no-interaction model is quite common, and easy to talk about.

The "random effect" for subjects is a little piece of random error, characteristic of an individual. We think of it as random because the individual was randomly sampled from a population. If, theoretically, the only reason that the measurements from a case are correlated is that each one is affected by this same little piece of under-performance or over-performance, the univariate approach represents a very good model.

The "random effect for a subject" idea implies a variance-covariance matrix of the dependent variables (say  $Y_1, \dots, 4$ ) with a *compound symmetry* structure.

$$\Sigma = \begin{bmatrix} \sigma^2 + \sigma_1 & \sigma_1 & \sigma_1 & \sigma_1 \\ \sigma_1 & \sigma^2 + \sigma_1 & \sigma_1 & \sigma_1 \\ \sigma_1 & \sigma_1 & \sigma^2 + \sigma_1 & \sigma_1 \\ \sigma_1 & \sigma_1 & \sigma_1 & \sigma^2 + \sigma_1 \end{bmatrix}.$$

Actually, compound symmetry is sufficient but not necessary for the univariate repeated F tests to be valid. All that's necessary is *sphericity*, which means the covariances of all differences among  $Y$ 's within a case are the same.

The classical univariate approach does have some weak points.

- The model is good if the *only* reason for correlation among the repeated measures is that one little piece of individuality added to each measurement by a subject. However, if there are other sources of covariation among the repeated measures (like learning, or fatigue, or memory of past performance), there is too much chance rejection of the null hypothesis. In this case the multivariate approach, with its unknown variance-covariance matrix, is more conservative. It is also more appropriate, if you have sufficient sample size.
- Even more conservative (overly so, if the assumptions of the multivariate approach are met) is the Greenhouse-Geisser correction, which compensates for the problem by reducing the error degrees of freedom.
- If the design is unbalanced (non-proportional  $n$ 's), the " $F$ -tests" of the classical univariate approach do not have an  $F$  distribution (even if all the statistical assumptions are satisfied), and it is unclear what they mean, if anything.
- Like the multivariate approach, the univariate approach to repeated measures analysis throws out a case if any of the observations are missing. (Did somebody say "mean substitution?" Oh no!)
- The univariate approach has real trouble with unequally spaced observations, and with very natural and high quality data sets where there may be different numbers of observations are collected for each individual.

#### **5.2.4 The covariance structure approach to repeated measures**

In the covariance structure approach, the data are set up to be read in a univariate manner, and one of the variables is a case identification, which will be used to determine which observations of a variable come from the

same case. Naturally, data lines from the same case should be adjacent in the file.

Instead of assuming independence or inducing compound symmetry within subjects by random effects assumptions, *one directly specifies the structure of the covariance matrix of the observations that come from the same subject.*

The following present no problem at all:

- Time-varying covariates (categorical, too)
- Unbalanced designs
- Unequally spaced observations
- Missing or unequal numbers of observations within subjects <sup>1</sup>
- More variables than subjects (but not more parameters than subjects)

It's implemented with SAS `proc mixed`. Only SAS seems to have it, though this should change as other software companies work to catch up.

- The “mixed” in `proc mixed` refers to mixed-model ANOVA. SAS `proc mixed` is indeed very strong in this, but we're just using it here for within-cases ANOVA. A good number of other powerful features will not be discussed here.
- Lots of different covariance structures are possible, including compound symmetry and unknown.
- Everything's still assumed multivariate normal.

```
/****** noise96c.sas *****/
options linesize=79 pagesize=250;
title 'Repeated measures on Noise data: Cov Struct Approach';
proc format;      value sexfmt    0 = 'Male'  1 = 'Female' ;

data loud;
  infile 'noise.dat'; /* Univariate data read */
```

---

<sup>1</sup>Provided this is unrelated to the variable being repeatedly measured. Like if the dependent variable is how sick a person is, and the data might be missing because the person is too sick to be tested, there is a serious problem.

```

input ident interest sex age noise time discrim ;
format sex sexfmt.;
label interest = 'Interest in topic (politics)'
      time      = 'Order of presenting noise level';

proc mixed method = ml;
  class age sex noise;
  model discrim = age|sex|noise;
  repeated / type = un subject = ident r;
  lsmeans age noise;

proc mixed method = ml;
  class age sex noise;
  model discrim = age|sex|noise;
  repeated / type = cs subject = ident r;

```

Now part of noise96c.lst.

#### The MIXED Procedure

##### Class Level Information

Class	Levels	Values
AGE	3	1 2 3
SEX	2	Female Male
NOISE	5	1 2 3 4 5

##### ML Estimation Iteration History

Iteration	Evaluations	Objective	Criterion
0	1	1521.4783527	
1	1	1453.7299937	0.00000000

Convergence criteria met.

R Matrix for Subject 1

Row	COL1	COL2	COL3	COL4	COL5
1	54.07988333	17.08300000	21.38658333	17.91785000	24.27668333
2	17.08300000	69.58763333	15.56748333	29.98861667	21.71448333
3	21.38658333	15.56748333	54.37978333	25.15906667	21.00126667
4	17.91785000	29.98861667	25.15906667	59.31531667	27.58265000
5	24.27668333	21.71448333	21.00126667	27.58265000	55.88941667

Covariance Parameter Estimates (MLE)

Cov Parm	Estimate	Std Error	Z	Pr >  Z
DIAG UN(1,1)	54.07988333	9.87359067	5.48	0.0001
UN(2,1)	17.08300000	8.22102992	2.08	0.0377
UN(2,2)	69.58763333	12.70490550	5.48	0.0001
UN(3,1)	21.38658333	7.52577602	2.84	0.0045
UN(3,2)	15.56748333	8.19197469	1.90	0.0574
UN(3,3)	54.37978333	9.92834467	5.48	0.0001
UN(4,1)	17.91785000	7.66900119	2.34	0.0195
UN(4,2)	29.98861667	9.15325956	3.28	0.0011
UN(4,3)	25.15906667	8.01928166	3.14	0.0017
UN(4,4)	59.31531667	10.82944565	5.48	0.0001
UN(5,1)	24.27668333	7.75870531	3.13	0.0018
UN(5,2)	21.71448333	8.52518917	2.55	0.0109
UN(5,3)	21.00126667	7.61610965	2.76	0.0058
UN(5,4)	27.58265000	8.24206793	3.35	0.0008
UN(5,5)	55.88941667	10.20396474	5.48	0.0001
Residual	1.00000000	.	.	.

Model Fitting Information for DISCRIM

Description	Value
Observations	300.0000
Variance Estimate	1.0000
Standard Deviation Estimate	1.0000
Log Likelihood	-1002.55
Akaike's Information Criterion	-1017.55
Schwarz's Bayesian Criterion	-1045.32
-2 Log Likelihood	2005.093
Null Model LRT Chi-Square	67.7484
Null Model LRT DF	14.0000
Null Model LRT P-Value	0.0000

Tests of Fixed Effects

Source	NDF	DDF	Type III F	Pr > F
AGE	2	54	5.95	0.0046
SEX	1	54	0.53	0.4716
AGE*SEX	2	54	0.41	0.6635
NOISE	4	216	18.07	0.0001
AGE*NOISE	8	216	1.34	0.2260
SEX*NOISE	4	216	0.99	0.4146
AGE*SEX*NOISE	8	216	1.30	0.2455

From the multivariate approach we had  $F = 5.35$ ,  $p < .001$  for age & approx  $F = 15.3$

Least Squares Means

Level	LSMEAN	Std Error	DDF	T	Pr >  T
AGE 1	38.66100000	1.21376060	54	31.85	0.0001
AGE 2	35.24200000	1.21376060	54	29.04	0.0001
AGE 3	32.76700000	1.21376060	54	27.00	0.0001
NOISE 1	39.82166667	0.94938474	216	41.94	0.0001
NOISE 2	36.83000000	1.07693727	216	34.20	0.0001
NOISE 3	35.30166667	0.95201351	216	37.08	0.0001
NOISE 4	34.38500000	0.99427793	216	34.58	0.0001
NOISE 5	31.44500000	0.96513744	216	32.58	0.0001

Now for the second mixed run we get the same kind of beginning, and then for compo

#### Tests of Fixed Effects

Source	NDF	DDF	Type III F	Pr > F
AGE	2	54	5.95	0.0046
SEX	1	54	0.53	0.4716
AGE*SEX	2	54	0.41	0.6635
NOISE	4	216	15.69	0.0001
AGE*NOISE	8	216	1.15	0.3338
SEX*NOISE	4	216	0.98	0.4215
AGE*SEX*NOISE	8	216	1.18	0.3096

Now proc glm will allow easy examination of residuals no matter which approach you take to repeated measures, provided the data are read in a univariate manner.

```

/***** noise96d.sas *****/
options linesize=79 pagesize=60;
title 'Repeated measures on Noise data: Residuals etc.';
proc format;      value sexfmt    0 = 'Male'  1 = 'Female' ;

data loud;
  infile 'noise.dat'; /* Univariate data read */
  input ident interest sex age noise time discrim ;
  format sex sexfmt.;

```



```

label interest = 'Interest in topic (politics)'
      time      = 'Order of presenting noise level';


proc glm;
  class age sex noise;
  model discrim = age|sex|noise;
  output out=resdata predicted=predis residual=resdis;


/* Look at some residuals */
proc sort; by time;
proc univariate plot;
  var resdis; by time;
proc plot;
  plot resdis * (ident interest);

/* Include time */
proc mixed method = ml;
  class age sex noise time;
  model discrim = time age|sex|noise;
  repeated / type = un subject = ident r;
  lsmeans time age noise;

```

Then I generated residuals from this new model using glm, and plotted again. Nothing.





Unfortunately time = 5 wound up on a separate page. When time is included in the model, the results get stronger but conclusions don't change.

Tests of Fixed Effects

Source	NDF	DDF	Type III F	Pr > F
TIME	4	266	17.67	0.0001
AGE	2	266	18.45	0.0001
SEX	1	266	1.63	0.2027
AGE*SEX	2	266	1.28	0.2789
NOISE	4	266	10.95	0.0001
AGE*NOISE	8	266	0.51	0.8488
SEX*NOISE	4	266	0.44	0.7784
AGE*SEX*NOISE	8	266	0.74	0.6573

Least Squares Means

Level	LSMEAN	Std Error	DDF	T	Pr >  T
TIME 1	29.54468242	0.91811749	266	32.18	0.0001
TIME 2	34.61557451	0.91794760	266	37.71	0.0001
TIME 3	36.18863723	0.92819179	266	38.99	0.0001
TIME 4	39.72344496	0.91838886	266	43.25	0.0001
TIME 5	37.71099421	0.93376736	266	40.39	0.0001
AGE 1	38.66100000	0.68895774	266	56.12	0.0001
AGE 2	35.24200000	0.68895774	266	51.15	0.0001
AGE 3	32.76700000	0.68895774	266	47.56	0.0001
NOISE 1	39.69226830	0.89132757	266	44.53	0.0001
NOISE 2	36.80608879	0.89274775	266	41.23	0.0001
NOISE 3	35.35302821	0.89130480	266	39.66	0.0001
NOISE 4	34.12899017	0.89502919	266	38.13	0.0001
NOISE 5	31.80295787	0.89180628	266	35.66	0.0001

Some nice covariance structures are available in `proc mixed`.

$$\text{Variance Components: type = vc} \quad \Sigma = \begin{bmatrix} \sigma_1^2 & 0 & 0 & 0 \\ 0 & \sigma_2^2 & 0 & 0 \\ 0 & 0 & \sigma_3^2 & 0 \\ 0 & 0 & 0 & \sigma_4^2 \end{bmatrix}$$

$$\text{Compound Symmetry: type = cs} \quad \Sigma = \begin{bmatrix} \sigma^2 + \sigma_1 & \sigma_1 & \sigma_1 & \sigma_1 \\ \sigma_1 & \sigma^2 + \sigma_1 & \sigma_1 & \sigma_1 \\ \sigma_1 & \sigma_1 & \sigma^2 + \sigma_1 & \sigma_1 \\ \sigma_1 & \sigma_1 & \sigma_1 & \sigma^2 + \sigma_1 \end{bmatrix}$$

$$\text{Unknown: type = un} \quad \Sigma = \begin{bmatrix} \sigma_1^2 & \sigma_{1,2} & \sigma_{1,3} & \sigma_{1,4} \\ \sigma_{1,2} & \sigma_2^2 & \sigma_{2,3} & \sigma_{2,4} \\ \sigma_{1,3} & \sigma_{2,3} & \sigma_3^2 & \sigma_{3,4} \\ \sigma_{1,4} & \sigma_{2,4} & \sigma_{3,4} & \sigma_4^2 \end{bmatrix}$$

$$\text{Banded: type = un(1)} \quad \Sigma = \begin{bmatrix} \sigma_1^2 & \sigma_5 & 0 & 0 \\ \sigma_5 & \sigma_2^2 & \sigma_6 & 0 \\ 0 & \sigma_6 & \sigma_3^2 & \sigma_7 \\ 0 & 0 & \sigma_7 & \sigma_4^2 \end{bmatrix}$$

$$\text{First order autoregressive: type = ar(1)} \quad \Sigma = \sigma^2 \begin{bmatrix} 1 & \rho & \rho^2 & \rho^3 \\ \rho & 1 & \rho & \rho^2 \\ \rho^2 & \rho & 1 & \rho \\ \rho^3 & \rho^2 & \rho & 1 \end{bmatrix}$$

There are more, including Toeplitz, Banded Toeplitz, Factor analytic, ARMA, and Spatial (covariance is a function of Euclidean distance between observations).