

Ordinary Multiple Regression with R*

```
> kars = read.table("http://www.utstat.toronto.edu/~brunner/appliedf12/data/mcars4.data")
> kars[1:4,]
  Cntry lper100k weight length
1    US     19.8   2178   5.92
2  Japan     9.9   1026   4.32
3    US    10.8   1188   4.27
4    US    12.5   1444   5.11
>
> attach(kars) # Variables are now available by name
> n = length(length); n
[1] 100
> # Make indicator dummy variables for Cntry
> # U.S. will be the reference category
> c1 = numeric(n); c1[Cntry=='Europ'] = 1
> table(c1,Cntry)
  Cntry
c1  Europ Japan US
  0     0    13 73
  1    14     0  0
> c2 = numeric(n); c2[Cntry=='Japan'] = 1
> table(c2,Cntry)
  Cntry
c2  Europ Japan US
  0    14     0 73
  1     0    13  0
>
> # Take a look at mean fuel consumption per country
> aggregate(lper100k,by=list(Cntry),FUN=mean)
Group.1      x
1  Europ 10.17857
2  Japan 10.68462
3    US 12.96438
> # Must specify a LIST of grouping factors
```

On average, the U.S. cars seem to be using more fuel. Back it up with a hypothesis test.

* Copyright information is on the last page.

Origin	c1	c2	$E(Y X=x) = \beta_0 + \beta_1 C_1 + \beta_2 C_2$
Europe	1	0	$\beta_0 + \beta_1$
Japan	0	1	$\beta_0 + \beta_2$
U.S.	0	0	β_0

```
> # One-factor ANOVA to compare means
> justcountry = lm(lper100k ~ c1+c2)
> summary(justcountry)
```

```
Call:
lm(formula = lper100k ~ c1 + c2)
```

```
Residuals:
    Min       1Q   Median       3Q      Max
-5.0644 -2.1644 -0.4644  2.5154  6.8356
```

```
Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  12.9644     0.3651  35.511 < 2e-16 ***
c1           -2.7858     0.9101  -3.061  0.00285 **
c2           -2.2798     0.9390  -2.428  0.01703 *
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 3.119 on 97 degrees of freedom
Multiple R-squared: 0.1203, Adjusted R-squared: 0.1022
F-statistic: 6.634 on 2 and 97 DF, p-value: 0.001993
```

```

>
> # Get nicer-looking ANOVA summary table
> is.factor(Cntry)
[1] TRUE
> jc2 = aov(lper100k~Cntry); summary(jc2) # aov is a wrapper for lm
      Df Sum Sq Mean Sq F value Pr(>F)
Cntry    2  129.1   64.55   6.634 0.00199 **
Residuals 97  943.8    9.73
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

>
> # Which means are different?
> TukeyHSD(jc2,ordered=T)
  Tukey multiple comparisons of means
    95% family-wise confidence level
    factor levels have been ordered

Fit: aov(formula = lper100k ~ Cntry)

$Cntry
      diff      lwr      upr      p adj
Japan-Europ 0.506044 -2.35364917 3.365737 0.9069443
US-Europ    2.785812  0.61956789 4.952056 0.0079628
US-Japan    2.279768  0.04470727 4.514829 0.0445191

>
> # The factor Cntry has dummy vars built in.
> # What are they?
> contrasts(Cntry) # Note alphabetical order
      Japan US
Europ    0  0
Japan    1  0
US       0  1

>
> summary(lm(lper100k~Cntry))

```

```
> summary(lm(lper100k~Cntry))
```

```
Call:
lm(formula = lper100k ~ Cntry)
```

```
Residuals:
    Min       1Q   Median       3Q      Max
-5.0644 -2.1644 -0.4644  2.5154  6.8356
```

```
Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  10.1786     0.8337  12.209 < 2e-16 ***
CntryJapan    0.5060     1.2014   0.421  0.67454
CntryUS       2.7858     0.9101   3.061  0.00285 **
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 3.119 on 97 degrees of freedom
Multiple R-squared: 0.1203, Adjusted R-squared: 0.1022
F-statistic: 6.634 on 2 and 97 DF, p-value: 0.001993
```

```
> # You can select the dummy variable coding scheme.
```

```
> contr.sum(3) # Effect coding
```

```
  [,1] [,2]
1     1     0
2     0     1
3    -1    -1
```

```
> contr.treatment(3,base=2) # Category 2 is the reference category
```

```
  1 3
1 1 0
2 0 0
3 0 1
```

```
> # U.S. as reference category again
> Country = Cntry
> contrasts(Country) = contr.treatment(3,base=3)
> summary(lm(lper100k~Country))
```

```
Call:
lm(formula = lper100k ~ Country)
```

```
Residuals:
    Min       1Q   Median       3Q      Max
-5.0644 -2.1644 -0.4644  2.5154  6.8356
```

```
Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  12.9644     0.3651   35.511 < 2e-16 ***
Country1     -2.7858     0.9101   -3.061  0.00285 **
Country2     -2.2798     0.9390   -2.428  0.01703 *
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 3.119 on 97 degrees of freedom
Multiple R-squared: 0.1203, Adjusted R-squared: 0.1022
F-statistic: 6.634 on 2 and 97 DF, p-value: 0.001993
```

Include covariates

Origin	c1	c2	$E(Y X=x) = \beta_0 + \beta_1X_1 + \beta_2X_2 + \beta_3C_1 + \beta_4C_2$
Europe	1	0	$(\beta_0 + \beta_3) + \beta_1X_1 + \beta_2X_2$
Japan	0	1	$(\beta_0 + \beta_4) + \beta_1X_1 + \beta_2X_2$
U.S.	0	0	$\beta_0 + \beta_1X_1 + \beta_2X_2$

```
> # Include covariates
> fullmodel = lm(lper100k ~ weight+length+Country)
> summary(fullmodel) # Look carefully at the signs!
```

Call:
`lm(formula = lper100k ~ weight + length + Country)`

Residuals:

Min	1Q	Median	3Q	Max
-4.5063	-0.8813	0.0147	1.3043	2.9432

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-7.276937	3.006354	-2.421	0.017399 *
weight	0.005457	0.001472	3.707	0.000352 ***
length	2.345968	0.980329	2.393	0.018676 *
Country1	1.487722	0.575633	2.584	0.011274 *
Country2	1.994239	0.584995	3.409	0.000958 ***

 Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.703 on 95 degrees of freedom
 Multiple R-squared: 0.7431, Adjusted R-squared: 0.7323
 F-statistic: 68.71 on 4 and 95 DF, p-value: < 2.2e-16

```
> # Test car size controlling for country
> anova(justcountry,fullmodel) # Full vs reduced
Analysis of Variance Table
```

Model 1: `lper100k ~ c1 + c2`
 Model 2: `lper100k ~ weight + length + Country`

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	97	943.81				
2	95	275.61	2	668.2	115.16	< 2.2e-16 ***

 Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
 > # I advise using anova ONLY to compare full and reduced models

```
>
> # Might as well test country controlling for size too.
> justsize = lm(lper100k ~ weight+length); summary(justsize)
```

```
Call:
lm(formula = lper100k ~ weight + length)
```

```
Residuals:
    Min     1Q  Median     3Q     Max
-4.3857 -1.0684 -0.0556  1.3077  4.0429
```

```
Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept) -3.617472   2.958472  -1.223  0.22439
weight       0.004949   0.001546   3.202  0.00185 **
length       1.835625   1.017349   1.804  0.07428 .
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 1.804 on 97 degrees of freedom
Multiple R-squared: 0.7058, Adjusted R-squared: 0.6997
F-statistic: 116.4 on 2 and 97 DF, p-value: < 2.2e-16
```

```
>
> anova(justsize,fullmodel)
Analysis of Variance Table
```

```
Model 1: lper100k ~ weight + length
Model 2: lper100k ~ weight + length + Country
  Res.Df  RSS Df Sum of Sq    F  Pr(>F)
1     97 315.64
2     95 275.61  2    40.035 6.8999 0.001592 **
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
>
```

```

> ##### Predictions and prediction intervals #####
>
> # Predict litres per 100 km for a Japanese car weighing 1295kg, 4.52m
long
> # (1990 Toyota Camry)
> betahat = fullmodel$coefficients; betahat
  (Intercept)      weight      length  Country1  Country2
-7.276936526  0.005456609  2.345968436  1.487721833  1.994238863
> contrasts(Country)
      1 2
Europ 1 0
Japan 0 1
US    0 0
> x1 = c(1,1295,4.52,0,1)
> sum(x1*betahat)
[1] 12.38739
>
> # Use the predict function
> # help(predict.lm)
>
> camry1990 = data.frame(weight=1295,length=4.52,Country='Japan');
camry1990
  weight length Country
1  1295   4.52   Japan
> predict(fullmodel,newdata=camry1990)
      1
12.38739
> # With 95 percent prediction interval (default)
> predict(fullmodel,newdata=camry1990, interval='prediction')
      fit      lwr      upr
1 12.38739  8.856608 15.91817

```



```

>
> # Multiple predictions
> cadillac1990 = data.frame(weight=1800,length=5.22,Country='US')
> volvo1990 = data.frame(weight=1371,length=4.823,Country='Europ')
> newcars = rbind(camry1990,cadillac1990,volvo1990); newcars
  weight length Country
1   1295  4.520   Japan
2   1800  5.220     US
3   1371  4.823   Europ
> is.data.frame(newcars)
[1] TRUE
> predict(fullmodel,newdata=newcars, interval='prediction')
      fit      lwr      upr
1 12.38739  8.856608 15.91817
2 14.79091 11.354379 18.22745
3 13.00640  9.481598 16.53121
>
> # It's not a bad idea to "predict" the observed data
> # Just look at the first 10 rows, for example
> cbind(lper100k,predict(fullmodel,interval='prediction'))[1:10,]
  lper100k      fit      lwr      upr
1    19.8 18.495691 15.012790 21.97859
2     9.9 10.450367  6.940724 13.96001
3    10.8  9.222800  5.747978 12.69762
4    12.5 12.590305  9.162660 16.01795
5    12.5 12.626349  9.219284 16.03341
6    12.5 12.626349  9.219284 16.03341
7    10.4  9.766491  6.236956 13.29603
8    13.2 14.570386 11.135730 18.00504
9    17.0 14.056832 10.515850 17.59782
10     9.2  8.330626  4.870298 11.79095

```

Warning message:

```

In predict.lm(fullmodel, interval = "prediction") :
  Predictions on current data refer to _future_ responses

```

This handout was prepared by Jerry Brunner, Department of Statistics, University of Toronto. It is licensed under a Creative Commons Attribution - ShareAlike 3.0 Unported License. Use any part of it as you like and share the result freely. The OpenOffice.org document is available from the course website:

<http://www.utstat.toronto.edu/~brunner/oldclass/appliedf12>