Supplementary Appendix

This appendix has been provided by the authors to give readers additional information about their work.

Supplement to: Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. N Engl J Med 2020;383:2603-15. DOI: 10.1056/NEJMoa2034577

(PDF last updated on December 17, 2020.)

SUPPLEMENTARY APPENDIX

List of Investigators	2
SARS-CoV-2 Testing Information	6
Table S1 Explanation of the Changes in Denominator Numbers in Various Analyses.	_
Table 31 Explanation of the Changes in Denominator Numbers in Various Analyses.	/
Table S2 Baseline Comorbidities	8
Table S3 Participants Reporting at Least 1 Adverse Event from Dose 1 (All Enrolled Participants)	9
Table S4 Vaccine Efficacy from 7 Days after Dose 2 by Underlying Comorbidities among Participants without	
Evidence of Infection Prior to 7 Days after Dose 2.	16
Evidence of fillection Frior to / Days after Dose 2.	.10
Table S5 Vaccine Efficacy of Severe COVID-19 Occurrence after Dose 1 (Modified Intention-to-Treat)	.11

List of Investigators

Name	Institute	Location	
Aberg, Judith	g, Judith Icahn School of Medicine at Mount Sinai		
Addo, Marylyn	Universitätsklinikum Hamburg-Eppendorf	Hamburg, Germany	
Akhan, Sıla	Kocaeli University Med	Kocaeli, Turkey	
Albertson, Timothy	University of California, Davis	Sacramento, CA, USA	
Al-Ibrahim, Mohamed	SNBL Clinical Pharmacology Center	Baltimore, MD, USA	
Altın, Sedat	Yedikule Chest Diseases and Thoracic Surgery Training and Research Hospital	Istanbul, Turkey	
Anderson, Corey	Clinical Research Consortium	Tempe, AZ, USA	
Andrews, Charles	Diagnostics Research Group	San Antonio, TX, USA	
Arora, Samir	Aventiv Research	Columbus, OH, USA	
Balik, Ismail	Ankara University Medical School	Ankara, Turkey	
Barnett, Elizabeth	Boston Medical Center	Boston, MA, USA	
Bauer, George	Benchmark Research	Metairie, LA, USA	
Baumann-Noss, Sybille	CRS Clinical Research Services Berlin	Berlin, Germany	
Berhe, Mezgebe	North Texas Infectious Diseases Consultants	Dallas, TX, USA	
Bradley, Paul	Meridian Clinical Research	Savannah, GA, USA	
Brandon, Donald	California Research Foundation	San Diego, CA, USA	
Brune, Daniel	Optimal Research	Peoria, IL, USA	
Burgher, Abram	The Pain Center of Arizona	Peoria, AZ, USA	
Butcher, Bain	Sterling Research Group	Cincinnati, OH, USA	
Butuk, David	Solaris Clinical Research	Meridian, ID, USA	
Cannon, Kevin	PMG Research of Wilmington	Wilmington, NC, USA	
Cardona, Jose	Indago Research and Health Center	Hialeah, FL, USA	
Cavelli, Rachael	State University of New York Upstate Medical University	Syracuse, NY, USA	
Chalhoub, Fadi	CNS Healthcare	Jacksonville, FL, USA	
Christensen, Shane	J. Lewis Research	Salt Lake City, UT, USA	
Christensen, Tom	Waterway Family Medicine	Little River, SC, USA	
Chu, Laurence	Benchmark Research	Austin, TX, USA	
Cox, Steven	Lynn Institute of Norman	Norman, OK, USA	
Crook, Gretchen	Austin Regional Clinic: ARC Wilson Parke	Austin, TX, USA	
Davis, Matthew	Rochester Clinical Research	Rochester, NY, USA	
Davit, Rajesh	Sterling Research Group	Cincinnati, OH, USA	
Denham, Douglas	Clinical Trials of Texas (CTT)	San Antonio, TX, USA	
Dever, Michael	Clinical Neuroscience Solutions	Orlando, FL, USA	
Donskey, Curtis	Louis Stokes Cleveland VA Medical Center	Cleveland, OH, USA	
Doust, Matthew	Hope Research Institute	Phoenix, AZ, USA	
Dunn, Michael	Quality Clinical Research	Omaha, NE, USA	
Earl, John	PMG Research of Hickory	Hickory, NC, USA	
Eder, Frank	United Medical Associates - Meridian Clinical Research	Binghamton, NY, USA	
Eich, Andreas	IKF Pneumologie	Frankfurt, Germany	
Ensz, David	Meridian Clinical Research	Dakota Dunes, SD, USA	
Essink, Brandon	Meridian Clinical Research	Omaha, NE, USA	
Falcone, Robert	Amici Clinical Research	Raritan, NJ, USA	
Falsey, Ann	University of Rochester/Rochester General Hospital	Rochester, NY, USA	
Farrington, Cecil	PMG Research of Salisbury	Salisbury, NC, USA	
armgon, Ccon	1 1910 Nescarcii di Sansuury	Sansoury, INC, USA	

Finberg, Robert University of Massachusetts Medical School Worcester, MA, USA Finn, Daniel Kentucky Pediatric/Adult Research Bardstown, KY, USA Fitz-Patrick, David East-West Medical Research Institute Honolulu, HI, USA Fortmann, Stephen Kaiser Permanente Center for Health Research Portland, OR, USA Thabazimbi, South Africa Fouche, Leon Tambotie Medical Centre Fragoso, Veronica Texas Center for Drug Development Houston, TX, USA Frenck, Robert Cincinnati Children's Hospital Cincinnati, OH, USA Fried, David Warwick, RI, USA Omega Medical Research Ventavia Research Group Keller, TX, USA Fuller, Gregory Fussell, Suzanne Long Beach Clinical Trials Services Long Beach, CA, USA Garcia-Diaz, Julia Ochsner Health New Orleans, LA, USA Gentry, Andrew Bozeman Health Clinical Research Bozeman, MT, USA Glover, Richard Alliance for Multispecialty Research Newton, KS, USA Greenbaum, Carla Benaroya Research Institute at Virginia Mason Seattle, WA, USA Grubb, Stephen Main Street Physicians Little River, SC, USA Hammitt, Laura Johns Hopkins Center for American Indian Health Whiteriver, AZ, USA Johns Hopkins Center for American Indian Health Chinle, AZ, USA Johns Hopkins Center for American Indian Health Shiprock, NM, USA Gallup, NM, USA Johns Hopkins Center for American Indian Health Harper, Charles Meridian Clinical Research Norfolk, NE, USA Wake Research Harper, Wayne Raleigh, NC, USA Hartman, Aaron Virginia Research Center Midlothian, VA, USA Heller, Robert Bayview Research Group Valley Village, CA, USA Hendrix, Ernest North Alabama Research Center Athens, AL, USA Herrington, Darrell Benchmark Research San Angelo, TX, USA Clinical Research Professionals Chesterfield, MO, USA Jennings, Timothy Sakarya University Faculty of Medicine Karabay, Oğuz Sakarya, Turkey Confluence Health/Wenatchee Valley Hospital & Clinics Kaster, Steven Wenatchee, WA, USA Katzman, Steven Michigan Center for Medical Research Farmington Hills, MI, USA Kingsley, Jeffrey Columbus Regional Research Institute Columbus, GA, USA Klein, Nicola Kaiser Permanente Santa Clara, CA, USA Kaiser Permanente Oakland, CA, USA Klein, Tracy Heartland Research Associates Wichita, KS, USA Koch, Mark Ventavia Research Group Fort Worth, TX, USA Köksal, Iftahar Acibadem University Hospital Atakent Istanbul, Turkey Koren, Michael Jacksonville Center for Clinical Research Jacksonville, FL, USA Kutner, Mark Miami, FL, USA Suncoast Research Group Lee, Marcus Trinity Clinical Research Tullahoma, TN, USA Huntingdon Park, CA, USA Leibowitz, Mark National Research Institute Levin, Michael Clinical Research Center of Nevada Las Vegas, NV, USA Fundacion INFANT Libster, Romina Buenos Aires, Argentina Fargo, ND, USA Lillestol, Michael Lillestol Research Somers Point, NJ, USA Lucasti, Christopher South Jersey Infectious Disease Studienzentrum Brinkum Dr. Lars Pohlmeier und Torsten Luttermann, Matthias Stuhr, Germany Drescher Manning, Mary Beth Velocity Clinical Research Cleveland, OH, USA Martin, Earl Martin Diagnostic Clinic Tomball, TX, USA MedPharmics Matherne, Paul Gulfport, MS, USA

McMurray, James Medical Affiliated Research Center (MARC) Huntsville, AL, USA

Mert, Ali Medipol University Hospital Istanbul, Turkey

Middleton, Randle Optimal Research Huntsville, AL, USA

Mitha, Essack Newtown Clinical Research Johannesburg, South Africa

Morawski, Emily Holston Medical Group Kingsport, TN, USA

Moreira, Edson Associação Obras Sociais Irmã Dulce and Oswaldo Cruz Bahia, Brazil

Foundation

Lynn Institute of Denver

Odekirk, Larry

Schultz, Armin

Mulligan, MarkNew York University Langone Vaccine CenterNew York, NY USAMurray, AlexanderPharmQuestGreensboro, NC, USAMussaji, MurtazaLinQ ResearchHouston, TX, USAMusungaie, DanyJongaie ResearchPretoria, South AfricaNell, HayleneTiervlei Trial Centre, Karl Bremer HospitalCape Town, South Africa

Ogbuagu, Onyema Yale University School of Medicine New Haven, CT, USA
Paolino, Kristopher State University of New York, Upstate Medical University Syracuse, NY, USA
Patel, Suchet Regional Clinical Research Endwell, NY, USA
Peterson, James J. Lewis Research Salt Lake City, UT, USA

Denver, CO, USA

Mannheim, Germany

Pickrell, Paul Tekton Research Austin, TX, USA

Polack, Fernando Fundacion INFANT Buenos Aires, Argentina

Poretz, Donald Clinical Alliance for Research and Education Annandale, VA, USA

Pod Course PMC Proceeds of Checkets

Raad, George PMG Research of Charlotte Charlotte, NC, USA Randall, William Dayton, OH, USA PriMed Clinical Research Rankin, Bruce Accel Research Sites - DeLand Clinical Research Unit DeLand, FL, USA Reynolds, Steven Collaborative Neuroscience Network Long Beach, CA, USA Atlanta Center for Medical Research Riesenberg, Robert Atlanta, GA, USA Acevedo Clinical Research Associates Miami, FL, USA Rodriguez, Hector

Rosen, Jeffrey Clinical Research of South Florida Coral Gables, FL, USA Rubino, John Raleigh Medical Group, PA Raleigh, NC, USA Rupp, Richard University of Texas Galveston, TX, USA Saiger, Salma Research Across America Mesquite, TX, USA Salata, Robert University Hospitals Cleveland Medical Center Cleveland, OH, USA Saleh, Jamshid Northern California Neurological Surgery Redding, CA, USA Schaefer, Axel Medizentrum Essen Borbeck Essen, Germany Schear, Martin Dayton Clinical Research Dayton, OH, USA

Schwartz, Howard Research Centers of America Hollywood, FL, USA Segall, Nathan Clinical Research Atlanta Stockbridge, GA, USA Seger, William HealthFirst Medical Group Fort Worth, TX, USA Senders, Shelly Senders Pediatrics South Euclid, OH, USA Sharp, Stephan Clinical Research Associates Nashville, TN, USA Cary, NC, USA Shoffner, Sylvia PMG Research of Cary

Simsek Yavuz, Serap İstanbul Faculty of Medicine, İstanbul University Istanbul, Turkey

CRS Clinical Research Services Mannheim

Sligh, TeresaProvidence Clinical ResearchNorth Hollywood, CA, USASmith, WilliamNew Orleans Center for Clinical ResearchNew Orleans, LA, USAStacey, HelenDiablo Clinical ResearchWalnut Creek, CA, USAStephens, MichaelFleming Island Center for Clinical ResearchFleming Island, FL, USA

Studdard, Harry Coastal Clinical Research Mobile, AL, USA
Tabak, Fehmi Istanbul University Cerrahpaşa Medical School Istanbul, Turkey

Talaat, Kawsar	Johns Hopkins School of Medicine	Baltimore, MD, USA
Thomas, Stephen	State University of New York, Upstate Medical University	Syracuse, NY, USA
Towner, William	Kaiser Permanente	Los Angeles, CA, USA
Tran, Van	Ventavia Research Group	Houston, TX, USA
Ünal, Serhat	Hacettepe University	Ankara, Turkey
Usdan, Lisa	CNS Healthcare	Memphis, TN, USA
Vanchiere, John	Louisiana State University Health Shreveport	Shreveport, LA, USA
Varano, Susann	Clinical Research Consulting	Milford, CT, USA
Wadsworth, L. Tyler	Sundance Clinical Research	St Louis, MO, USA
Walsh, Edward	University of Rochester/Rochester General Hospital	Rochester, NY, USA
Walter, Emmanuel	Duke Human Vaccine Institute	Durham, NC, USA
Wappner, Diego	Fundacion INFANT	Buenos Aires, Argentina
Whiles, Rick	Holston Medical Group	Bristol, TN, USA
Williams, Hayes	Achieve Clinical Research	Birmingham, AL, USA
Wilson, Jonathan	Piedmont Medical Research of Winston-Salem	Winston-Salem, NC, USA
Winkle, Peter	Anaheim Clinical Trials	Anaheim, CA, USA
Winokur, Patricia	University of Iowa	Iowa City, IA, USA
Wolf, Thomas	Prairie Fields Family Medicine	Fremont, NE, USA
Yozviak, Joseph	Lehigh Valley Hospital Cedar Crest	Allentown, PA, USA
Zerbini, Cristiano	Centro Paulista de Investigação Clínica - CEPIC	Sao Paolo, Brazil

SARS-CoV-2 Testing Information

Testing for SARS-CoV-2 virus was conducted using the Cepheid Xpert Xpress SARS-CoV-2 PCR test.

Testing for SARS-CoV-2 antibodies was conducting using the Roche Elecsys $^{\otimes}$ Anti-SARS-CoV-2 antibody test.

Figure/Table Number	Figure/Table Title	Population(s)/Sample Size	Explanation
Figure 1	Disposition of participants (CONSORT)	All enrolled population N=37,706 "main safety subset"	All randomized ≥16 years of age, N=43,548
			• [minus 99 non-vaccinated, 1 no ICD]
			Vaccinated N=43,448
			Main safety subset (N=37,706) needed to have been enrolled by October 9, 2020 for EUA application
Figure 2	Local and Systemic Reactions Reported within	Reactogenicity subset of ≥16 years old	Per protocol
	7 Days after Receipt of 30 µg BNT162b2 or Placebo by Age Group	N=8,183	
Figure 3	Efficacy of BNT162b2 against COVID-19	N=43,355 (modified intention-to-treat)	All randomized >=12 years of age N= 43,651
	Occurrence after Dose 1		• [minus 99 non-vaccinated, 1 no ICD]
			Vaccinated (dose 1 efficacy) N=43,551
			• [minus 196 HIV+]
			All efficacy N=43,355
Table 1	Demographics	N=37,706 main safety subset	As above
Table 2	Vaccine Efficacy against	1st primary efficacy endpoint:	Evaluable population:
	COVID-19 from 7 Days after Dose 2 [Primary	Includes those without evidence of prior infection (N=36,523)	 received 2 vaccinations as randomized
	Endpoints]	and maintage affice as an amaint.	 no major protocol
		2nd primary efficacy endpoint: Includes those with and without evidence of prior infection (N=40,137)	deviations Excludes HIV+
Table 3	Vaccine Efficacy Overall and by Subgroup in Participants without Evidence of Infection Prior to 7 Days After Dose 2	N=36,523 (same as 1st primary endpoint)	
Table S2	Baseline Comorbidities	N=37,706 main safety subset	
Table S3	Participants Reporting at Least 1 Adverse Event From Dose 1 (All Enrolled Participants)	N=43,252	Vaccinated N=43,448 minus 196 HIV+
Table S4	Vaccine Efficacy from 7 Days After Dose 2 by Underlying Comorbidities among Participants without Evidence of Infection Prior to 7 Days after Dose 2	N=36,523 (same as 1st primary endpoint)	
Table S5	Vaccine Efficacy of Severe COVID-19 Occurrence after Dose 1 (Modified Intention-to-Treat)	N=43,355 (modified intention-to-treat)	See comments to Figure 3

Table S1 | Explanation of the Changes in Denominator Numbers in Various Analyses.

	BNT162b2 (30 μg) (N ^a =18860)	Placebo (Na=18846)	Total (Na=37706)
Charlson Comorbidity Index Category	n ^b (%)	n ^b (%)	n^b (%)
Participants with any Charlson comorbidity	3934 (20.9)	3809 (20.2)	7743 (20.5)
AIDS/HIV	59 (0.3)	62 (0.3)	121 (0.3)
Any malignancy	733 (3.9)	662 (3.5)	1395 (3.7)
Cerebrovascular disease	195 (1.0)	166 (0.9)	361 (1.0)
Chronic pulmonary disease	1478 (7.8)	1453 (7.7)	2931 (7.8)
Congestive heart failure	88 (0.5)	83 (0.4)	171 (0.5)
Dementia	7 (0.0)	11 (0.1)	18 (0.0)
Diabetes with chronic complication	99 (0.5)	113 (0.6)	212 (0.6)
Diabetes without chronic complication	1473 (7.8)	1478 (7.8)	2951 (7.8)
Hemiplegia or paraplegia	13 (0.1)	21 (0.1)	34 (0.1)
Leukemia	12 (0.1)	10 (0.1)	22 (0.1)
Lymphoma	22 (0.1)	32 (0.2)	54 (0.1)
Metastatic solid tumor	4 (0.0)	3 (0.0)	7 (0.0)
Mild liver disease	125 (0.7)	89 (0.5)	214 (0.6)
Moderate or severe liver disease	1 (0.0)	2 (0.0)	3 (0.0)
Myocardial infarction	194 (1.0)	188 (1.0)	382 (1.0)
Peptic ulcer disease	52 (0.3)	71 (0.4)	123 (0.3)
Peripheral vascular disease	124 (0.7)	117 (0.6)	241 (0.6)
Renal disease	123 (0.7)	133 (0.7)	256 (0.7)
Rheumatic disease	62 (0.3)	56 (0.3)	118 (0.3)

Table S2 | **Baseline Comorbidities.** Baseline comorbid conditions are classified according to the Charlson Comorbidity Index (Charlson M, TP Szatrowski, J Peterson, J. Gold. Validation of a combined comorbidity index. J Clin Epidemiol 1994; 47:1245-51.). a. N = number of participants in the specified group. This value is the denominator for the percentage calculations. b. n = Number of participants with the specified characteristic. Participants with multiple occurrences within each category are counted only once. For 'Participants with any Charlson comorbidity', n = number of participants reporting at least 1 occurrence of any Charlson comorbidity.

	BNT162b2 (30 μg) (N ^a =21621)	Placebo (Na=21631)
Adverse Event	n ^b (%)	n ^b (%)
Any event	5770 (26.7)	2638 (12.2)
Related ^c	4484 (20.7)	1095 (5.1)
Severe	240 (1.1)	139 (0.6)
Life-threatening	21 (0.1)	24 (0.1)
Any serious adverse event	126 (0.6)	111 (0.5)
Related ^c	4 (0.0)	0
Severe	71 (0.3)	68 (0.3)
Life-threatening	21 (0.1)	23 (0.1)
Any adverse event leading to withdrawal	37 (0.2)	30 (0.1)
Related ^c	16 (0.1)	9 (0.0)
Severe	13 (0.1)	9 (0.0)
Life-threatening	3 (0.0)	6 (0.0)
Death	2 (0.0)	4 (0.0)

Table S3 | **Participants Reporting at Least 1 Adverse Event from Dose 1 (All Enrolled Participants).** The 'all enrolled' population included all participants who received at least 1 dose of vaccine irrespective of follow-up time. a. N = number of participants in the specified group. This value is the denominator for the percentage calculations. b. n = Number of participants reporting at least 1 occurrence of the specified event category. For 'any event', n = the number of participants reporting at least 1 occurrence of any event. c. Assessed by the investigator as related to investigational product.

		BNT162b2 (30 μg) (N ^a =18198)		Placebo (Na=18325)		
Efficacy Endpoint Subgroup	n1 ^b	Surveillance $Time^{c} (n2^{d})$	n1 ^b	Surveillance $Time^{c}$ $(n2^{d})$	VE (%)	(95% CI ^e)
Overall	8	2.214 (17411)	162	2.222 (17511)	95.0	(90.0, 97.9)
At risk ^f						
Yes	4	1.025 (8030)	86	1.025 (8029)	95.3	(87.7, 98.8)
No	4	1.189 (9381)	76	1.197 (9482)	94.7	(85.9, 98.6)
Age group (years) and at risk						
16-64 and not at risk	4	0.962 (7671)	69	0.964 (7701)	94.2	(84.4, 98.5)
16–64 and at risk	3	0.744 (5878)	74	0.746 (5917)	95.9	(87.6, 99.2)
≥65 and not at risk	0	0.227 (1701)	7	0.233 (1771)	100.0	(29.0, 100.0)
≥65 and at risk	1	0.281 (2147)	12	0.279 (2109)	91.7	(44.2, 99.8)
Obese ^g						
Yes	3	0.763 (6000)	67	0.782 (6103)	95.4	(86.0, 99.1)
No	5	1.451 (11406)	95	1.439 (11404)	94.8	(87.4, 98.3)
Age group (years) and obese						
16-64 and not obese	4	1.107 (8811)	83	1.101 (8825)	95.2	(87.3, 98.7)
16-64 and obese	3	0.598 (4734)	60	0.609 (4789)	94.9	(84.4, 99.0)
≥65 and not obese	1	0.343 (2582)	12	0.338 (2567)	91.8	(44.5, 99.8)
≥65 and obese	0	0.165 (1265)	7	0.173 (1313)	100.0	(27.1, 100.0

Table S4 | Vaccine Efficacy from 7 Days after Dose 2 by Underlying Comorbidities among Participants without Evidence of Infection Prior to 7 Days after Dose 2. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period. a. N = 1000 number of participants in the specified group. b. 1000 number of participants meeting the endpoint definition. c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period. d. 1000 number of participants at risk for the endpoint. e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time. f. At risk is defined as having at least one of the Charlson Comorbidity Index categories or obesity (body mass index [BMI] ≥30 kg/m²). g. Obese is defined as BMI ≥30 kg/m².

	BNT162b2 (30 μg) (Na=21669)			Placebo (Na=21686)		
Efficacy Endpoint Subgroup	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance $Time^{c}\ (n2^{d})$	VE (%)	(95% CI ^e)
Severe COVID-19 occurrence after Dose 1	1	4.021 (21314)	9	4.006 (21259)	88.9	(20.1, 99.7)
After Dose 1 to before Dose 2	0		4		100.0	(-51.5, 100.0)
Dose 2 to 7 days after Dose 2	0		1		100.0	(-3800.0, 100.0)
≥7 Days after Dose 2	1		4		75.0	(-152.6, 99.5)

Table S5 | Vaccine Efficacy of Severe COVID-19 Occurrence after Dose 1 (Modified Intention-to-Treat). a. N = number of participants in the specified group. b. n1 = Number of participants meeting the endpoint definition. c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from Dose 1 to the end of the surveillance period. d. n2 = Number of participants at risk for the endpoint. e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method (adjusted for surveillance time for overall row).