

Regression diagnostics with R*

```
> sat = read.table("http://www.utstat.utoronto.ca/~brunner/data/legal/openSAT.data.txt")
> head(sat)
  VERBAL MATH GPA
1     623  509 2.6
2     454  471 2.3
3     643  700 2.4
4     585  719 3.0
5     719  710 3.1
6     693  643 2.9
> mod1 = lm(GPA ~ VERBAL+MATH, data=sat); summary(mod1)
```

Call:

```
lm(formula = GPA ~ VERBAL + MATH, data = sat)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.24875	-0.35113	0.04659	0.38745	1.03527

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.6062975	0.4414062	1.374	0.171
VERBAL	0.0023072	0.0005522	4.178	4.42e-05 ***
MATH	0.0009999	0.0006093	1.641	0.102


Signif. codes: 0 ‘***’ 0.001 ‘**’ 0.01 ‘*’ 0.05 ‘.’ 0.1 ‘ ’ 1

Residual standard error: 0.5484 on 197 degrees of freedom
Multiple R-squared: 0.1161, Adjusted R-squared: 0.1071
F-statistic: 12.93 on 2 and 197 DF, p-value: 5.284e-06

```
> attach(sat) # Make variable names accessible
>
```


* Copyright information is on the last page.

```
> # Plot y-hat versus y  
> yhat = mod1$fitted.values  
> plot(yhat,GPA)  
>
```




```
> cor(GPA,yhat)^2 # r-squared - R-squared  
[1] 0.1160179
```

```
> # Plot y-hat versus residuals  
> e = mod1$residuals  
> plot(yhat,e)
```



```
> cor(yhat,e) # Zero  
[1] 2.898153e-16
```

```
> # Compare plot of standardized residuals  
> sr = rstandard(mod1)  
> plot(yhat,sr,ylab='Standardized Residual')
```




```

> # Three look like possible outliers: Investigate
> id = 1:200
> suspect = id[sr < -2.5]
> cbind(sat[suspect,],yhat[suspect],sr[suspect])
   VERBAL MATH  GPA yhat[suspect] sr[suspect]
20     497  640 0.69      2.392965 -3.128203
51     645  548 1.16      2.642644 -2.735853
73     516  569 0.94      2.365985 -2.620870

> # Studentized deleted residuals are t-statistics
> sdr = rstudent(mod1) # Studentized deleted residuals
> # Bonferroni critical value for n=200 tests, at joint alpha = 0.05 level
> dfe = mod1$df.residual; dfe
[1] 197
> alpha = 0.05; a = alpha/200; bcrit = qt(1-a/2,dfe-1); bcrit
[1] 3.730706
> summary(sdr)
   Min. 1st Qu. Median      Mean 3rd Qu.      Max.
-3.201000 -0.672800  0.048340 -0.001731  0.710900  2.317000


```

```
> hist(sdr,xlab='Studentized Deleted Residual',
+ main='Histogram of Studentized Deleted Residuals')
```



```
> # Should we worry about normality? Not if maximum diagonal element  
of the hat matrix is less than 0.2  
> hii = hatvalues(mod1)  
> summary(hii) # Max should be less than 0.2 for robustness.  
Min. 1st Qu. Median Mean 3rd Qu. Max.  
0.005071 0.007824 0.011310 0.015000 0.019100 0.062370  
> hist(hii)
```

Histogram of h_{ii}



Trees Data

```
> rm(list=ls()) # Remove everything to start
> head(trees)
   Girth Height Volume
1    8.3     70   10.3
2    8.6     65   10.3
3    8.8     63   10.2
4   10.5     72   16.4
5   10.7     81   18.8
6   10.8     83   19.7
> attach(trees)
> mod1 = lm(Volume ~ Girth + Height)
> summary(mod1)


Call:
lm(formula = Volume ~ Girth + Height)

Residuals:
    Min      1Q  Median      3Q     Max 
-6.4065 -2.6493 -0.2876  2.2003  8.4847 


Coefficients:
            Estimate Std. Error t value Pr(>|t|)    
(Intercept) -57.9877    8.6382  -6.713 2.75e-07 ***
Girth        4.7082    0.2643  17.816 < 2e-16 ***
Height       0.3393    0.1302   2.607   0.0145 *  
---
Signif. codes:  0 '****' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 3.882 on 28 degrees of freedom
Multiple R-squared:  0.948, Adjusted R-squared:  0.9442 
F-statistic: 255 on 2 and 28 DF,  p-value: < 2.2e-16

>
> plot(yhat,Volume)
```



```
> rstud1 = rstudent(mod1) # Studentized deleted residuals  
> plot(Girth,rstud1)
```



```

> plot(Height,rstud1) # No pattern
> Girthsq = Girth^2 # Polynomial term -- literally square it
> mod2 = lm(Volume ~ Girth + Girthsq + Height); summary(mod2)

Call:
lm(formula = Volume ~ Girth + Girthsq + Height)

Residuals:
    Min      1Q  Median      3Q     Max 
-4.2928 -1.6693 -0.1018  1.7851  4.3489 

Coefficients:
            Estimate Std. Error t value Pr(>|t|)    
(Intercept) -9.92041   10.07911 -0.984 0.333729    
Girth        -2.88508    1.30985 -2.203 0.036343 *  
Girthsq       0.26862    0.04590  5.852 3.13e-06 *** 
Height        0.37639    0.08823  4.266 0.000218 *** 
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.625 on 27 degrees of freedom
Multiple R-squared:  0.9771, Adjusted R-squared:  0.9745 
F-statistic: 383.2 on 3 and 27 DF,  p-value: < 2.2e-16

> rstud2 = rstudent(mod2)
> plot(Girth,rstud2) # Nothing
> plot(Height,rstud2) # Nothing
>
> # Look at prediction intervals
> cbind(Volume[1:5],predict(mod1,interval='predict')[1:5,])
   fit      lwr      upr
1 10.3  4.837660 -3.561809 13.23713
2 10.3  4.553852 -3.962908 13.07061
3 10.2  4.816981 -3.809144 13.44311
4 16.4 15.874115  7.690594 24.05764
5 18.8 19.869008 11.451358 28.28666

Warning message:
In predict.lm(mod1, interval = "predict") :
  predictions on current data refer to _future_ responses

>
> cbind(Volume[1:5],predict(mod2,interval='predict')[1:5,])
   fit      lwr      upr
1 10.3 10.985950  4.902269 17.06963
2 10.3  9.600406  3.566753 15.63406
3 10.2  9.205421  3.163767 15.24708
4 16.4 16.501775 10.954762 22.04879
5 18.8 20.451204 14.746331 26.15608

Warning message:
In predict.lm(mod2, interval = "predict") :
  predictions on current data refer to _future_ responses

```

This document is licensed under a Creative Commons Attribution - ShareAlike 3.0 Unported License:
http://creativecommons.org/licenses/by-sa/3.0/deed.en_us. Use any part of it as you like
and share the result freely.