

Sample Questions: Maximum Likelihood Part 1

STA312 Spring 2019. Copyright information is at the end of the last page.

Let X_1, \dots, X_n be independent Pareto random variables with density $f(x|\theta) = \begin{cases} \frac{\theta}{x^{\theta+1}} & \text{for } x \geq 1 \\ 0 & \text{for } x < 1 \end{cases}$

where $\theta > 0$. The Pareto has a decreasing density with a heavy right tail, sometimes used as a model for the unequal distribution of wealth.

1. Derive a formula for the maximum likelihood estimate of θ . Include the second derivative test. Show your work and **circle your final answer**.

2. Give a formula for \hat{v}_n , the estimated asymptotic variance of $\hat{\theta}_n$. Show a little work.

3. The file <http://www.utstat.toronto.edu/~brunner/data/legal/pareto.data.txt> has a set of raw data. Calculate

- (a) The maximum likelihood estimate $\hat{\theta}_n$.
- (b) A 95% confidence interval for θ .

Your answers are numbers. Circle and label them. Bring your printout to the quiz.

```
> rm(list=ls())
> x = scan("http://www.utstat.toronto.edu/~brunner/data/legal/pareto.data.txt")
Read 150 items
> x
 [1] 5.47 2.54 4.01 1.22 2.74 4.99 1.24 4.35 227.65
[10] 3.20 4.35 1.02 1.17 3.49 1.61 10.43 9.04 1.07
[19] 4.80 1.14 1.41 36.62 5.38 1.98 1.43 1.54 1.42
[28] 1.06 1.68 1.44 1.52 2.25 1.62 1.01 53.79 1.11
[37] 1.52 28.39 15.55 3.96 2.73 6.43 4.35 1.29 2.04
[46] 1.04 1.68 1.89 1.78 2.57 1.39 5.49 1.07 1.74
[55] 5.68 1.43 1.58 42.42 2.11 1.07 1.27 1.03 1.02
[64] 10.92 1.43 2.18 6.28 1.81 4.42 1.93 2.39 3.75
[73] 1.65 1.01 1.31 2.66 1.08 1.36 1.22 2.20 2.79
[82] 1.11 2.01 3.11 2.02 2.21 1.05 5.69 1.16 5.47
[91] 2.19 1.22 1.37 1.37 1.63 3.55 1.13 1.26 1.21
[100] 1.35 4.36 4.59 1.47 2.22 4.34 2.19 1.80 1.68
[109] 31.22 3.63 1.01 1.60 2.39 2.21 1.22 1.54 2.39
[118] 1008.06 3.37 1.30 4.46 1.01 7.51 1.14 1.02 1.12
[127] 2.83 1.79 1.67 1.45 1.74 1.66 1.01 1.34 1.12
[136] 1.30 1.18 10.96 2.28 2.33 1.51 1.41 1.19 1.40
[145] 1.83 1.33 31.98 5.86 3.28 1.53
> thetahat = 1/mean(log(x)); thetahat
[1] 1.100816
> n = length(x); vhat = thetahat^2/n; se = sqrt(vhat); se
[1] 0.08988125
> low95 = thetahat - 1.96*se; up95 = thetahat + 1.96*se
> c(low95,up95) # 95% CI
[1] 0.9246488 1.2769833
```

4. For the Pareto distribution, the well-known 80-20 rule (80% of the wealth is held by 20% of the population) corresponds to a value of $\theta = 1.16$. Using a two-sided large-sample Z -test and the usual $\alpha = 0.05$ significance level, test whether these data are compatible with $\theta = 1.16$.
- (a) There are two critical values, one for the lower tail and one for the upper tail. What are they? The answers are numbers.
 - (b) What is the value of the test statistic? The answer is a number. Circle it.
 - (c) Use R to calculate the 2-sided p -value. The answer is a number.

```
> # Critical value(s). Just say plus and minus 1.96, or ...
> c(qnorm(0.025),qnorm(0.975))
[1] -1.959964  1.959964
>
> Z = (thetahat-1.16)/se; Z
[1] -0.6584683
>
> pvalue = 2 * (1-pnorm(abs(Z))); pvalue
[1] 0.5102373
```

- (d) Do you reject the null hypothesis? Answer Yes or No.
- (e) Are the results statistically significant? Answer Yes or No.
- (f) Do these data contradict claim that $\theta = 1.16$? Answer Yes or No.

This assignment was prepared by [Jerry Brunner](#), Department of Mathematical and Computational Sciences, University of Toronto. It is licensed under a [Creative Commons Attribution - ShareAlike 3.0 Unported License](#). Use any part of it as you like and share the result freely. The \LaTeX source code is available from the course website:

<http://www.utstat.toronto.edu/~brunner/oldclass/312s19>