

# Random Vectors<sup>1</sup>

STA442/2101 Fall 2016

---

<sup>1</sup>See last slide for copyright information.

## Background Reading: Renscher and Schaalje's *Linear models in statistics*

- Chapter 3 on Random Vectors and Matrices

## Random Vectors and Matrices

A *random matrix* is just a matrix of random variables. Their joint probability distribution is the distribution of the random matrix. Random matrices with just one column (say,  $p \times 1$ ) may be called *random vectors*.

The expected value of a matrix is defined as the matrix of expected values. Denoting the  $p \times c$  random matrix  $\mathbf{X}$  by  $[X_{i,j}]$ ,

$$E(\mathbf{X}) = [E(X_{i,j})].$$

Immediately we have natural properties like

$$\begin{aligned} E(\mathbf{X} + \mathbf{Y}) &= E([X_{i,j} + Y_{i,j}]) \\ &= [E(X_{i,j} + Y_{i,j})] \\ &= [E(X_{i,j}) + E(Y_{i,j})] \\ &= [E(X_{i,j})] + [E(Y_{i,j})] \\ &= E(\mathbf{X}) + E(\mathbf{Y}). \end{aligned}$$

## Moving a constant through the expected value sign

Let  $\mathbf{A} = [a_{i,j}]$  be an  $r \times p$  matrix of constants, while  $\mathbf{X}$  is still a  $p \times c$  random matrix. Then

$$\begin{aligned} E(\mathbf{AX}) &= E\left(\left[\sum_{k=1}^p a_{i,k} X_{k,j}\right]\right) \\ &= \left[E\left(\sum_{k=1}^p a_{i,k} X_{k,j}\right)\right] \\ &= \left[\sum_{k=1}^p a_{i,k} E(X_{k,j})\right] \\ &= \mathbf{A}E(\mathbf{X}). \end{aligned}$$

Similar calculations yield  $E(\mathbf{AXB}) = \mathbf{A}E(\mathbf{X})\mathbf{B}$ .

## Variance-Covariance Matrices

Let  $\mathbf{X}$  be a  $p \times 1$  random vector with  $E(\mathbf{X}) = \boldsymbol{\mu}$ . The *variance-covariance matrix* of  $\mathbf{X}$  (sometimes just called the *covariance matrix*), denoted by  $cov(\mathbf{X})$ , is defined as

$$cov(\mathbf{X}) = E \left\{ (\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top \right\}.$$

$$\text{cov}(\mathbf{X}) = E \left\{ (\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top \right\}$$

$$\begin{aligned} \text{cov}(\mathbf{X}) &= E \left\{ \begin{pmatrix} X_1 - \mu_1 \\ X_2 - \mu_2 \\ X_3 - \mu_3 \end{pmatrix} \begin{pmatrix} X_1 - \mu_1 & X_2 - \mu_2 & X_3 - \mu_3 \end{pmatrix} \right\} \\ &= E \left\{ \begin{pmatrix} (X_1 - \mu_1)^2 & (X_1 - \mu_1)(X_2 - \mu_2) & (X_1 - \mu_1)(X_3 - \mu_3) \\ (X_2 - \mu_2)(X_1 - \mu_1) & (X_2 - \mu_2)^2 & (X_2 - \mu_2)(X_3 - \mu_3) \\ (X_3 - \mu_3)(X_1 - \mu_1) & (X_3 - \mu_3)(X_2 - \mu_2) & (X_3 - \mu_3)^2 \end{pmatrix} \right\} \\ &= \begin{pmatrix} E\{(X_1 - \mu_1)^2\} & E\{(X_1 - \mu_1)(X_2 - \mu_2)\} & E\{(X_1 - \mu_1)(X_3 - \mu_3)\} \\ E\{(X_2 - \mu_2)(X_1 - \mu_1)\} & E\{(X_2 - \mu_2)^2\} & E\{(X_2 - \mu_2)(X_3 - \mu_3)\} \\ E\{(X_3 - \mu_3)(X_1 - \mu_1)\} & E\{(X_3 - \mu_3)(X_2 - \mu_2)\} & E\{(X_3 - \mu_3)^2\} \end{pmatrix} \\ &= \begin{pmatrix} \text{Var}(X_1) & \text{Cov}(X_1, X_2) & \text{Cov}(X_1, X_3) \\ \text{Cov}(X_1, X_2) & \text{Var}(X_2) & \text{Cov}(X_2, X_3) \\ \text{Cov}(X_1, X_3) & \text{Cov}(X_2, X_3) & \text{Var}(X_3) \end{pmatrix}. \end{aligned}$$

So, the covariance matrix  $\text{cov}(\mathbf{X})$  is a  $p \times p$  symmetric matrix with variances on the main diagonal and covariances on the off-diagonals.



## Matrix of covariances between two random vectors

Let  $\mathbf{X}$  be a  $p \times 1$  random vector with  $E(\mathbf{X}) = \boldsymbol{\mu}_x$  and let  $\mathbf{Y}$  be a  $q \times 1$  random vector with  $E(\mathbf{Y}) = \boldsymbol{\mu}_y$ . The  $p \times q$  matrix of covariances between the elements of  $\mathbf{X}$  and the elements of  $\mathbf{Y}$  is

$$C(\mathbf{X}, \mathbf{Y}) = E \left\{ (\mathbf{X} - \boldsymbol{\mu}_x)(\mathbf{Y} - \boldsymbol{\mu}_y)^\top \right\}.$$

# Adding a constant has no effect

On variances and covariances

- $cov(\mathbf{X} + \mathbf{a}) = cov(\mathbf{X})$
- $C(\mathbf{X} + \mathbf{a}, \mathbf{Y} + \mathbf{b}) = C(\mathbf{X}, \mathbf{Y})$

It's clear from the definitions:

- $cov(\mathbf{X}) = E \{ (\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top \}$
- $C(\mathbf{X}, \mathbf{Y}) = E \{ (\mathbf{X} - \boldsymbol{\mu}_x)(\mathbf{Y} - \boldsymbol{\mu}_y)^\top \}$

So sometimes it is useful to let  $\mathbf{a} = -\boldsymbol{\mu}_x$  and  $\mathbf{b} = -\boldsymbol{\mu}_y$ .

## Analogous to $Var(aX) = a^2 Var(X)$

Let  $\mathbf{X}$  be a  $p \times 1$  random vector with  $E(\mathbf{X}) = \boldsymbol{\mu}$  and  $cov(\mathbf{X}) = \boldsymbol{\Sigma}$ , while  $\mathbf{A} = [a_{i,j}]$  is an  $r \times p$  matrix of constants. Then

$$\begin{aligned} cov(\mathbf{AX}) &= E \left\{ (\mathbf{AX} - \mathbf{A}\boldsymbol{\mu})(\mathbf{AX} - \mathbf{A}\boldsymbol{\mu})^\top \right\} \\ &= E \left\{ \mathbf{A}(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top \right\} \\ &= E \left\{ \mathbf{A}(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top \mathbf{A}^\top \right\} \\ &= \mathbf{A} E \left\{ (\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top \right\} \mathbf{A}^\top \\ &= \mathbf{A} cov(\mathbf{X}) \mathbf{A}^\top \\ &= \mathbf{A} \boldsymbol{\Sigma} \mathbf{A}^\top \end{aligned}$$

This slide show was prepared by **Jerry Brunner**, Department of Statistics, University of Toronto. It is licensed under a **Creative Commons Attribution - ShareAlike 3.0 Unported License**. Use any part of it as you like and share the result freely. The  $\text{\LaTeX}$  source code is available from the course website:  
<http://www.utstat.toronto.edu/~brunner/oldclass/appliedf16>